

AN EXAMPLE OF A BLOCH FUNCTION

RICHARD M. TIMONEY

Department of Mathematics
Indiana University
Bloomington, Indiana 47401 U.S.A.

(Received November 3, 1978)

ABSTRACT. A Bloch function is exhibited which has radial limits of modulus one almost everywhere but fails to belong to H^p , for each $0 < p \leq \infty$.

KEY WORDS AND PHRASES. Bloch function.

AMS (MOS) SUBJECT CLASSIFICATION (1970) CODES. 30A78

1. INTRODUCTION.

The purpose of this note is to give an example which seems to be useful in settling several questions about Bloch functions.

Let E be the subset of the complex plane \mathbb{C} consisting of the closed unit disc together with the Gaussian integers \mathbb{Z}^2 . Let G be the complement of E in \mathbb{C} . Let $g : D \rightarrow G$ be the analytic universal covering map of G given by the uniformization theorem (D denotes the unit disc).

PROPOSITION. The function g is an unbounded Bloch function with the properties

- (i) g has a radial limit $g(e^{i\theta})$ at almost every point $e^{i\theta}$ of the unit circle.

- (ii) the function $g(e^{i\theta})$ is of modulus one almost everywhere on the unit circle,
- (iii) g is the reciprocal of a singular inner function, and so g does not belong to any H^p class.

Bloch functions on the unit disc may be defined as those analytic functions f on D for which the radii of the schlicht discs in the range of f are bounded above. The Bloch functions are somewhat analogous to functions in the disc algebra--Bloch functions can be characterized (see [1]) as those analytic functions which are uniformly continuous when D is given the hyperbolic metric and \mathbb{C} the Euclidean metric. Since Bloch functions may be characterized (see [1]) as those analytic functions f on D for which the quantity $|f'(z)|(1 - |z|^2)$ is bounded for $z \in D$, it follows that the modulus of a Bloch function grows rather slowly--at most as fast as $\log(1/(1 - |z|))$. Because functions in the disc algebra and bounded functions have good boundary behaviour, it is natural to ask about boundary values of Bloch functions--in particular about radial boundary values. (It is shown in [4] that a Bloch function has a radial limit at a point of the unit circle if and only if it has a non-tangential limit there.)

In [5], Pommerenke gave an example of a Bloch function with radial limits almost nowhere. The example given here is constructed in a similar way, but it contrasts with Pommerenke's in that it shows that Bloch functions which have radial limits almost everywhere need not be particularly well-behaved.

The example answers a question posed by Joseph Cima (private communication). He asked whether a Bloch function which has radial limits

almost everywhere and has the additional property that the boundary function belongs to L^p need be in H^p . The function g provides a negative answer to this question since $g(e^{i\theta}) \in L^\infty$ while $g \notin H^p$ for any $0 < p \leq \infty$. In fact g does not belong to the class N^+ (see [2] p. 25) which contains H^p for every p .

PROOF. It is evident that g is an unbounded Bloch function. Also, to verify properties (i), (ii) and (iii), it is clearly sufficient to verify (iii).

To establish (iii), consider the analytic function $f = 1/g$ on D . The function f is bounded (by 1) and is the universal covering map: $D \rightarrow D - K$, where K is the countable set

$$\{0\} \cup \{1/(m+in) \mid m, n \in \mathbb{Z}, |m+in| > 1\}.$$

Being a bounded analytic function, f has radial limits almost everywhere on the unit circle. It is easy to see from the properties of covering maps that these radial limits are either of modulus 1 or else belong to K . To complete the proof that f is a singular inner function, it is only necessary to show that the radial limit $f(e^{i\theta})$ belongs to K on a subset of the unit circle of measure zero.

But, for each $k \in K$ it is true that the set of $e^{i\theta}$ for which $f(e^{i\theta}) = k$ has measure zero (see [2] p. 17). Since K is countable, it follows that the set of $e^{i\theta}$ for which $f(e^{i\theta})$ belongs to K also has measure zero. The proof is now complete.

The example may also be viewed as elucidating the almost total lack of relationships between the class \mathcal{B} of Bloch functions on D and the subclasses H^p and N^+ of the Nevanlinna class N (see [2]). The only

containment which holds between \mathcal{B} and the other classes is the relation $H^\infty \subseteq \mathcal{B}$. It is known that $H^p \not\subseteq \mathcal{B}$ for any $0 < p < \infty$ and that $\mathcal{B} \not\subseteq N$. The example g given above belongs to $\mathcal{B} \cap N$ but not to N^+ . The fact $\mathcal{B} \not\subseteq N$ is shown by the example of Pommerenke's [5] mentioned above.

Finally, the example given here can be modified to show that there is no $\delta > 0$ such that an analytic function $f : D \rightarrow \mathbb{C}$ satisfying

$$f(e^{i\theta}) = \lim_{r \rightarrow 1} f(re^{i\theta}) = 1$$

almost everywhere on the unit circle must have a disc of radius δ in its range. (Merely replace \mathbb{Z}^2 by $\delta\mathbb{Z}^2$ in the construction of g). This answers a question raised by J.S. Hwang. By contrast, he showed (see [3]) that a singular inner function (for example) must have a (Schlicht) disc of radius at least $2B/e$ in its range, where B denotes Bloch's constant.

REFERENCES

1. Anderson, J.M., J.G. Clunie, Ch. Pommerenke. On Bloch Functions and Normal Functions, J. Reine Angew. Math. 270(1974) 12-37.
2. Duren, P. H^p Spaces, Academic Press, 1970.
3. Hwang, J.S., On an Extremal Property of Doob's Class (preprint, 1978).
4. Lehto, O., K.I. Virtanen. Boundary behaviour and normal meromorphic functions. Acta Math. 97(1957) 47-65.
5. Ch. Pommerenke. On Bloch functions. J. London Math. Soc. (2) 2(1970) 689-695.

Special Issue on Abstract Differential and Difference Equations

Call for Papers

The purpose of this special issue is to highlight current progress in abstract differential and difference equations and their applications. Survey articles devoted to important problems in the field are welcome.

This special issue will focus on the following topics:

- Almost periodic and almost automorphic solutions
- Topological methods
- Equations with multivalued operators
- Infinite dimensional dynamical systems with discrete or continuous time
- Particular solutions such as standing and traveling waves
- Applications to mathematical physics, engineering, and biology

Articles published in this special issue will be granted 50% reduction on the applied Article Processing Charges.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/ade/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	February 1, 2010
First Round of Reviews	May 1, 2010
Publication Date	August 1, 2010

Lead Guest Editor

G. M. N'Guérékata, Department of Mathematics, Morgan State University, Baltimore, MD 21251, USA;
gaston.n'guerekata@morgan.edu

Guest Editors

T. Diagana, Department of Mathematics, Howard University, Washington, DC 20005, USA;
tdiagana@howard.edu

Special Issue on Models, Methods, and Applications of Dynamics and Control in Engineering Sciences: State of the Art

Call for Papers

Since the current literature on dynamics and control is scattered over a range of journals, books, chapters of books, and a large number of conference proceedings, which are often difficult to obtain, the goal of this special issue of MPE is to present papers, containing complete reviews on dynamics models, available in the current literature, to classify them, and to discuss their applications and limitations. In this special issue, the authors could recommend appropriate models and control criteria for various applications on engineering and sciences and suggest directions for further works. It is also open for critical reviews, open problems, and future developments. Discussions on MEMS and NEMS are also encouraged.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/mpe/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	August 1, 2009
First Round of Reviews	November 1, 2009
Publication Date	February 1, 2010

Lead Guest Editor

Jose Manoel Balthazar, Department of Statistics, Applied Mathematics and Computation, State University of São Paulo at Rio Claro, P.O. Box 178, Rio Claro 13500-230, SP, Brazil; jmbaltha@rc.unesp.br

Guest Editors

Paulo Batista Gonçalves, Civil Engineering Department, Pontifical Catholic University of Rio de Janeiro (PUC-Rio), 22453-900 Rio de Janeiro, RJ, Brazil; paulo@puc-rio.br

Stefano Lenzi, Istituto di Scienza e Tecnica delle Costruzioni, Università Politecnica delle Marche, via Brezze Bianche, Monte D'Ago, Ancona 60131, Italy; lenci@univpm.it

Yuri Vladimirovich Mikhlin, Department of Applied Mathematics, Kharkov Polytechnical Institute, National Technical University, Kharkov 61002, Ukraine; muv@kpi.kharkov.ua

Special Issue on Nonlinear Time Series: Computations and Applications

Call for Papers

Nonlinear time series attracts researchers from many areas of sciences and technologies, ranging from mathematics and physics to computer science. Two categories of nonlinear time series will be considered in this special issue. The one is fractal time series with power law; the other is chaotic series.

The focus of this special issue is on the computations of nonlinear time series toward the applications to various issues in science and engineering. It would be an opportunity of extending the research field of fractals, chaos, wavelets, applied mathematics, and applied statistics in all aspects of theoretical and practical study.

We are soliciting original high-quality research papers on topics of interest connected with the nonlinear time series that include, but are not limited to the following main topics:

- Fractal or chaotic time series, its modeling, and computations
- Dynamical systems relating to fractal or chaotic time series
- Wavelet analysis and its application to nonlinear time series

In addition, applications to pulses, the Internet, bio-engineering, industrial management science, cryptography and information security, control engineering, and complex networks are welcome.

Before submission authors should carefully read over the journal's Author Guidelines, which are located at <http://www.hindawi.com/journals/mpe/guidelines.html>. Prospective authors should submit an electronic copy of their complete manuscript through the journal Manuscript Tracking System at <http://mts.hindawi.com/> according to the following timetable:

Manuscript Due	February 1, 2010
First Round of Reviews	May 1, 2010
Publication Date	August 1, 2010

Lead Guest Editor

Ming Li, School of Information Science & Technology, East China Normal University, No. 500, Dong-Chuan Road, Shanghai 200241, China; mli@ee.ecnu.edu.cn

Guest Editors

Massimo Scalia, Dipartimento di Matematica, Sapienza Università di Roma, Roma, Italy; massimo.scalia@uniroma1.it

Cristian Toma, Faculty of Applied Sciences, Politehnica University of Bucharest, Hagi-Ghita 81, 060032 Bucharest, Romania; cgtoma@physics.pub.ro