

β -Neurexin Is a Ligand for the *Staphylococcus aureus* MSCRAMM SdrC

E. Magda Barbu^{1,2}, Vannakambadi K. Ganesh¹, Shivasankarappa Gurusiddappa¹, R. Chris Mackenzie², Timothy J. Foster³, Thomas C. Sudhof⁴, Magnus Höök^{1*}

1 Center for Infectious & Inflammatory Diseases, Institute of Biosciences and Technology, Texas A&M Health Science Center, Houston, Texas, United States of America, **2** Microbiology and Molecular Genetics Department, University of Texas Health Science Center at Houston, Houston, Texas, United States of America, **3** Moyné Institute of Preventive Medicine, Department of Microbiology, Trinity College, Dublin, Ireland, **4** Neuroscience Institute, Department of Molecular and Cellular Physiology, Stanford University, Palo Alto, California, United States of America

Abstract

Gram-positive bacteria contain a family of surface proteins that are covalently anchored to the cell wall of the organism. These cell-wall anchored (CWA) proteins appear to play key roles in the interactions between pathogenic organisms and the host. A subfamily of the CWA has a common structural organization with multiple domains adopting characteristic IgG-like folds. The identified microbial surface components recognizing adhesive matrix molecules (MSCRAMMs) belong to this subfamily, as does SdrC from *S. aureus*. However, an interactive host ligand for the putative MSCRAMM SdrC was not previously identified. We have screened a phage display peptide library and identified a peptide sequence found in β -neurexin that binds SdrC. A synthetic peptide corresponding to the identified sequence as well as a recombinant form of the β -neurexin 1 exodomain binds SdrC with high affinity and specificity. Furthermore, expression of SdrC on bacteria greatly enhances microbial adherence to cultured mammalian cells expressing β -neurexin on their surface. Taken together, our experimental results demonstrate that β -neurexin is a ligand for SdrC. This interaction involves a specific sequence located in the N-terminal region of the mammalian protein and the N₂N₃ domain of the MSCRAMM. The fact that these two proteins interact when expressed on the appropriate cells demonstrates the functionality of the interaction. Possible implications of this interaction are discussed.

Citation: Barbu EM, Ganesh VK, Gurusiddappa S, Mackenzie RC, Foster TJ, et al. (2010) β -Neurexin Is a Ligand for the *Staphylococcus aureus* MSCRAMM SdrC. PLoS Pathog 6(1): e1000726. doi:10.1371/journal.ppat.1000726

Editor: Ambrose Cheung, Dartmouth Medical School, United States of America

Received: June 1, 2009; **Accepted:** December 15, 2009; **Published:** January 15, 2010

Copyright: © 2010 Barbu et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Funding source: NIH/NIAMS AI20624-22 "Molecular Analysis of Staphylococcus aureus Host Interactions" (<http://www.niams.nih.gov/>). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: mhook@ibt.tamhsc.edu

Introduction

The Gram-positive opportunistic pathogen *Staphylococcus aureus* can cause a spectrum of infections in humans and animals that differ in severity. Some relatively minor skin infections, such as folliculitis, impetigo and cellulitis, can progress to life threatening diseases like sepsis, endocarditis, osteomyelitis and pneumonia [1].

The molecular pathogenic mechanisms of different staphylococcal infections are incompletely understood but studies suggested that a critical factor for the pathogenic success of this organism depends on its ability to adhere effectively to multiple host tissues [2,3,4]. The adhesins mediating staphylococcal adherence and colonization often target the extracellular matrix of the host and hence belong to the MSCRAMM family [4]. In Gram-positive bacteria, many MSCRAMMs are cell-wall-anchored (CWA) proteins [5,6] with a similar structural organization [7]. These proteins contain an amino terminal signal sequence followed by an A-region that often harbors the ligand-binding sites. The A-region is comprised of sub-domains (called N-domains) adopting an immunoglobulin G-like (IgG-like) fold [8]. Sometimes the A-region is followed by a B-region containing repeated β -sandwich modules of unknown function. In the case of the Sdr-subfamily of staphylococcal MSCRAMMs, the B-region is accompanied by a

repeat (R) domain composed of multiple Ser-Asp dipeptide repeats (SD-repeat or Sdr) (reviewed in [9]).

The ligand-binding activity of SdrG, a fibrinogen (Fg)-binding MSCRAMM of *Staphylococcus epidermidis* [10], was shown to proceed via a "dock, lock and latch" mechanism [7,11]. A crystal structure of a Fg-based peptide in complex with the SdrG_{N2N3} domain suggested that the peptide "docks" into the groove formed between the N2 and N3 domain. Upon binding the C-terminal extension of the N3 domain is redirected to cover and "lock" the ligand peptide in place and further stabilizes the complex by complementing a β -sheet in the N2 domain, thus functionally serving as a "latch" [11]. The back of the latching trench contains a motif, TYTFTDYVD, conserved in several other staphylococcal MSCRAMMs. Genome-based bioinformatics of five Gram-positive bacterial strains revealed that all organisms contain proteins with predicted IgG-like folded domains, an LPXTG-motif and a TYTFTDYVD-motif. Therefore, the "dock, lock and latch" mechanism was proposed as a common ligand binding strategy among these proteins.

In the current study, we used SdrC as bait for screening a phage library displaying 12-mer linear peptides (New England Biolabs Ph.D.TM-12 Phage Display Peptide Library). The *sdrC* gene is the first gene in a bi- or tri-partite gene cluster that also contains either *sdrD* or *sdrD* and *sdrE*. The predicted domain organization of SdrC

Author Summary

Staphylococcus aureus is an opportunistic pathogen, distinguished by its potential to cause serious and life-threatening infections in animals and humans. The ability of this bacterium to adhere to host tissues is considered an early, essential event in the disease process and contributes to the success of the organism as a pathogen. Adherence to host tissues is mediated by a subfamily of cell-wall anchored proteins named MSCRAMMs (microbial surface component recognizing adhesive matrix molecules). Work in our laboratory suggested that many of these proteins share a common ligand binding mechanism targeting linear amino acid sequences. Based on these observations, we hypothesized that screening a phage display library of random peptides may identify receptors for MSCRAMMs. Using this method, we demonstrate that the putative MSCRAMM SdrC recognizes a sequence in the neuronal protein β-neurexin. Furthermore, we show that intact β-neurexin 1 is a functional ligand for the *S. aureus* MSCRAMM SdrC. Successful implementation of this approach may open avenues for the identification of additional host ligands and the design of anti-staphylococcal therapeutics able to inhibit these interactions.

is similar to that of SdrG with the N2 and N3 sub-domains of the A-region adopting an IgG-like fold [12,13].

The phage display strategy, used in this study, identified beta (β) neurexin as a ligand for the orphan MSCRAMM SdrC. Subsequent biochemical and cell biology experiments confirmed that β-neurexin is a ligand for SdrC.

Neurexins are neuronal cell adhesion molecules that interact with neuroligins and appear to play an important role in synapse function (reviewed in [14]). Previously, the neurexins were discovered as the receptor for the neurotoxin α-latrotoxin of the black-widow spider [15].

Results

Validation of the phage display method to identify MSCRAMM ligands

Analysis of peptides or protein segments displaced on the surface of phage is a useful method to identify partners in protein-protein interaction systems [16,17,18,19]. To determine if this method can be used to identify ligands for MSCRAMMs from Gram-positive bacteria, we performed a validation experiment using SdrG_{N2N3} as bait in a screen of a commercially available 12-mer linear library (New England Biolabs Ph.D.-12 Phage Display Peptide Library). We carried out three successive rounds of panning against immobilized SdrG_{N2N3}. To confirm that the enrichment of the phage clones between panning rounds was the result of specific interactions, the phage pool before and after each step of amplification was also tested for binding to BSA. We found that the number of phage binding to SdrG increased relative to the number of phage binding to BSA (Table 1) suggesting a specific enrichment for SdrG. We then sequenced the DNA purified from ten phage plaques obtained after the third round of panning. Alignment of the inserted amino acid sequences revealed FSARG as a consensus in the majority of the clones (Table 2). This result was consistent with previous results from our laboratory identifying FFSARG as the binding motif in Fg for SdrG [7]. The successful identification of the SdrG binding sequence suggested that phage display of peptide libraries could be used to find binding partners for orphan MSCRAMMs.

Table 1. Number of transducing units (TUs) obtained after each panning against BSA and SdrG_{N2N3}.

Panning*	Ligand	Total bound TUs
1	BSA	1 × 10 ⁴
	SdrG	4 × 10 ⁵
2	BSA	1 × 10 ²
	SdrG	3 × 10 ⁷
3	BSA	10
	SdrG	2 × 10 ⁸

*Number of input phage was 1.5 × 10¹¹ for each panning.
doi:10.1371/journal.ppat.1000726.t001

Identification of the putative ligand-binding domain of SdrC

Sequence alignments between the A-region of SdrC (SdrC_A or SdrC₅₂₋₄₉₆) and previously identified staphylococcal MSCRAMMs revealed a modest identity (less than 20%). In contrast, a comparison of the predicted structure of SdrC_A with the determined structures of the crystallized staphylococcal MSCRAMMs ClfA and SdrG pointed to close structural similarities [7,20,21]. A subsegment of the SdrC A-region corresponding to residues 178–496 was predicted by PHYRE (<http://www.sbg.bio.ic.ac.uk/phyre/>) to adopt a structure similar to those determined for the N2N3 sub-domain of the Fg-binding MSCRAMMs ClfA (PHYRE $\epsilon = 3.1 \times 10^{-32}$) and SdrG (PHYRE $\epsilon = 9.6 \times 10^{-32}$). Similar to the N2N3 domains of ClfA and SdrG, SdrC₁₇₈₋₄₉₆ likely contains two IgG-like folded domains and has a TYTFIDYVD-like “latching cleft” motif in the predicted N2 sub-domain. Therefore, we designated SdrC₁₇₈₋₄₉₆ as the N2N3 domain and SdrC₅₂₋₁₇₇ as the N1 domain.

Both SdrC₅₂₋₄₉₆ (SdrC_A) and SdrC₁₇₈₋₄₉₆ (SdrC_{N2N3}) were expressed with an N-terminal His tag in *E. coli* (Fig. 1A). SDS-PAGE analysis of purified recombinant proteins indicated apparent molecular weights of 60 KDa (Fig. 1B) and 45 KDa, respectively (Fig. 1C). Matrix-assisted laser desorption ionization mass spectrometry suggested that the masses of the purified recombinant proteins are close to the theoretical molecular masses calculated from the primary amino acid sequences (51,194 KDa compared to 50,924 KDa for SdrC_A and 38,358 KDa compared

Table 2. Peptide motifs binding to SdrG_{N2N3}.

Peptide sequence
VYPTLH FS ARGSG
QSQWPVLF FS ARG
SNIPFQ FS ARGPG
SENLQ FS ARGPGG
HMEGSELS FS ARG
AHQDETLA FS ARG
NHPPFNALS FS SRG
YADLLAQ FS SKSG
YLPSQ IS AIGRAG
NPVIRYAS RS SHSG

*consensus sequence shown in bold.
doi:10.1371/journal.ppat.1000726.t002




Figure 1. Structural organization of SdrC and recombinant proteins. (A) Schematic representation of SdrC domain structure. S, signal sequence; A region underlined composed of N1, N2 and N3, B1 and B2, R repeats, R, serine-aspartic acid repeat region; W, wall-spanning fragment; M, transmembrane domain; C, cytoplasmic tail; LPETG, cell wall anchoring motif; VAAPQ, cleavage site; the arrow indicated the cleavage position. Also shown, are the His-tag recombinant SdrC proteins used in this study. (B), Coomassie-stained SDS-PAGE gel of pure recombinant SdrC_A. (C) Coomassie-stained SDS-PAGE gel of the recombinant SdrC_A after 2 weeks storage at 4°C. doi:10.1371/journal.ppat.1000726.g001

to 38,340 KDa for SdrC_{N2N3}). Thus, the recombinant SdrC proteins migrate aberrantly on SDS-PAGE. Aberrant migration on SDS-PAGE is common for the recombinant A-region of MSCRAMMs and thought to be due to their hydrophilic index [10,12,22]. The recombinant A-region was quickly degraded to a single proteolytically stable segment with an apparent molecular mass of 45 KDa. The N-terminal fragment released was apparently further degraded and was rarely detected on SDS-PAGE. N-terminal sequence analysis of the truncated, resistant segment revealed that cleavage had occurred between Ala¹⁷⁷ and Ala¹⁷⁸. Mass spectrometry indicated a molecular weight of 38,358 KDa compared with 38,340 KDa calculated from the predicted amino acid sequence of the stable fragment suggesting that the cleavage event occurred only at the N-terminus of the A-region. Cleavage of the N-terminal sub-segment (N1) of the A-region has also been observed with other recombinant MSCRAMMs (ClfA, ClfB, SdrG) and may be explained by disordered segments in the N1 sub-domains as well as the presence of minute amounts of contaminating proteases in the MSCRAMM preparations [7,23,24]. The proteolysis resistant fragment of SdrC_A corresponds to the tandem IgG-like domains (N2N3) as predicted by PHYRE (<http://www.sbg.bio.ic.ac.uk/phyre/>).

Identification of β -neurexin as a potential binding partner for SdrC_{N2N3} using phage display

We screened the New England Biolabs Ph.D.-12 Phage Display Peptide Library using immobilized recombinant SdrC_{N2N3} as bait. After the 3rd round of panning, fifty randomly picked plaques were selected and the corresponding phages were screened again for binding to SdrC_{N2N3}, to SdrG_{N2N3} and to BSA as controls. All clones bound to SdrC, however, phage from eight clones displayed significantly higher binding to SdrC_{N2N3} compared to SdrG_{N2N3} and BSA (Fig. 2A). The clone inserts were sequenced and an alignment of the deduced amino acid sequences is shown in

Table 3. From these sequences we could identify a degenerate consensus sequence (P,T,A)HH(I,M)HHFH(G,R,S,Q,T,A) which was then used for a pattern search of the human protein database employing an algorithm allowing for zero, one or two residue mismatches. Zero mismatches returned only neurexin 1 β , one mismatch returned neurexin 1 β and neurexin 2 β and two mismatches returned neurexin 1 β , neurexin 2 β , neurexin 3 β and a T-type voltage-dependent calcium channel (Table S1). All proteins returned by the search were screened for the presence of the consensus sequence within the extracellular domain of the protein. As shown in Fig. 2B, an alignment of the N-terminal extracellular segments of the β -neurexin isoforms revealed variations of the consensus sequence identified by phage display. In addition, the peptide sequence displayed by phage number one confirms a sequence found in neurexin 1 β and the sequenced displayed by phage eight is very similar to a sequence found in neurexin 3 β [25].

The amino-terminal segments of the β -neurexins contain an unusually long signal sequence (~50 aa), a neuroligin binding site and a glycosylation site in close proximity to the transmembrane domain. The C-terminal cytoplasmic segment contains a PDZ domain presumably involved in intracellular signaling [25]. The putative SdrC binding site identified by phage display corresponds to amino acid residues 10 to 16 and maps in the neuroligin-binding domain exposed on the cell surface.

Recombinant SdrC_{N2N3} binds recombinant neurexin 1 β (Nrx1 β)

The phage display library screening identified beta neurexins as potential binding partners for SdrC. To explore a possible interaction between the recombinant forms of SdrC_{N2N3} and the exodomain of Nrx1 β we expressed in *E. coli* two recombinant Nrx1 β fragments with a C-terminal GST tag (Fig. 3A). The fragment designated Nrx₄₇₋₂₅₅ corresponds to the first 208 amino acids residues of Nrx1 β protein which contains the putative




Figure 2. Identification of β-neurexins as the potential SdrC-binding partner. (A), Phage obtained after the third panning round bound specifically to SdrC_{N2N3}. Fifty random phage clones obtained after the 3rd round of panning were incubated with immobilized SdrC_{N2N3}, SdrG_{N2N3} and BSA. Shown are the eight phage clones with the highest binding affinity (*p<0.001) for SdrC_{N2N3} in comparison with SdrG_{N2N3}, BSA or fd-tet (insertless phage). (B), Consensus sequence search. Pattern searches with the degenerated consensus (P,T,A)HH(I,M)HHFH(G,R,S,Q,T,A) against a human protein database returned β-neurexin isoforms as SdrC-ligands. Consistent residues of the consensus are highlighted in bold, variable residues are highlighted in gray.
doi:10.1371/journal.ppat.1000726.g002

SdrC-binding site identified by phage display (amino acid residues 57 through 63). Nr_x₇₀₋₂₅₅ is a truncated version of the segment described above in which the putative SdrC-binding site was deleted (Fig. 3B). Both recombinant Nr_x1β-GST proteins were immobilized on 96-multiwell plates and used in a solid phase binding assay. SdrC_A bound the immobilized Nr_x₄₇₋₂₅₅ in a concentration dependent manner but did not bind to either Nr_x₇₀₋₂₅₅ or the purified fusion partner GST (Fig. 3C). Similar results were obtained when the intact SdrC_{N2N3} region (Fig. 3D) or the proteolytically resistant sub-segment released from the recombinant SdrC_A (data not shown) were used as probes.

To further demonstrate that HHHHFFH is the specific binding site in Nr_x1β for SdrC, we used a synthetic peptide (sp) with a sequence corresponding to the N-terminus of Nr_x1β, SLGAH-

HHHFFHGSSKHHS (spNr_x_{wt}), in attempts to inhibit the binding between SdrC_{N2N3} and Nr_x₄₇₋₂₅₅. As a control we used a peptide with the same residues but where the sequence was scrambled to HSHIKLHSHGHSFGHA (spNr_x_{scr}). Recombinant SdrC_{N2N3} (0.5 μM) was incubated with increasing concentration of either peptide prior to being added to the Nr_x coated plate. spNr_x_{wt} inhibited the interaction between SdrC_{N2N3} and Nr_x₄₇₋₂₅₅ in a concentration dependent manner. We did not observe an inhibition of the MSCRAMM-β-neurexin interaction when spNr_x_{scr} was used in the assay (Fig. 3E). The observed binding of recombinant SdrC to Nr_x1β and the subsequent synthetic peptide inhibition assay suggests that the interaction between these proteins is specific and that the sequence motif identified in the phage display experiment represents the ligand-binding site in Nr_x1β for SdrC.

Table 3. Peptide motifs binding to SdrC_{N2N3}.

	Peptide sequence*
Peptide 1	HWRT HHIHFFH QOG
Peptide 2	MSP HHMHHS HGHG
Peptide 3	AKLA HHIHFFH G
Peptide 4	WVP HHIHFFH RAG
Peptide 5	Y THHHHS WRLHTG
Peptide 6	A HHPH AWRHSHKG
Peptide 7	ST FHFH THKARHG
Peptide 8	HS QHRRF HHTYPG

*consensus sequence shown in bold.
doi:10.1371/journal.ppat.1000726.t003

SdrC binds the Nr_x_{wt} peptide with high affinity

We used fluorescence polarization to determine the dissociation constant for the SdrC_{N2N3}-Nr_x1β peptide interaction. spNr_x_{wt} and spNr_x_{scr} were labeled with fluorescein and incubated with increasing concentrations of recombinant MSCRAMM. SdrC_{N2N3} bound spNr_x_{wt} in a concentration dependent, saturable manner with a K_D of 2.5±0.5×10⁻⁷ M. We did not detect a significant binding of SdrC_{N2N3} to the fluorescein-labeled spNr_x_{scr}. A control MSCRAMM, ClfB_{N2N3}, did not bind to the fluorescein-labeled spNr_x_{wt} (Fig. 4A).

To eliminate the possibility that the binding was dependent of the fluorescein label introduced in the wild-type peptide, we tested the ability of unlabeled spNr_x_{wt} and spNr_x_{scr} to inhibit the binding of SdrC_{N2N3} to fluorescein-labeled spNr_x_{wt}. Increasing concentrations of unlabeled peptides were incubated with SdrC_{N2N3} for three hours at room temperature before the fluorescent polarization experiment




Figure 3. Structural organization of β -neurexins. (A) A cartoon representation of β -neurexin domain structure. S, signal sequence; NL, neuroigin-binding domain; Glyc, glycosylation domain; TM, transmembrane domain; PDZ, intracellular signaling domain. Also shown, are the recombinant proteins used in this study. (B) Coomassie-stained SDS-PAGE gel of GST-tagged neurexin 1 β domains recombinant proteins. Increasing concentrations of SdrC_A (C) and SdrC_{N2N3} (D) were incubated with immobilized Nrx_{47-255} (black square), Nrx_{70-255} (black triangle) or GST (black circle) for 1 hour at room temperature. The apparent K_D (concentrations required for half maximum binding) values were $1.85 \pm 0.94 \times 10^{-7}$ M for SdrC_A and $1.26 \pm 0.28 \times 10^{-7}$ M for SdrC_{N2N3}. (E) SdrC₁₇₈₋₄₉₆ (0.5 μM) was incubated with increasing concentrations of spNrx_{wt} peptide (black square) or spNrx_{scr} peptide (black triangle) at room temperature. After 2 hours, the mixture was incubated with immobilized Nrx_{47-255} . The values represented here are the mean \pm SD of triplicates from three experiments. doi:10.1371/journal.ppat.1000726.g003

was performed. Fig. 4B shows that the unlabeled spNrx_{wt} but not the spNrx_{scr} inhibited the binding between SdrC_{N2N3} and the fluorescein-labeled spNrx_{wt}. These results demonstrated that SdrC binds with relatively high affinity to the identified amino acid sequence motif in β -neurexin.

SdrC mediates bacterial attachment to cells expressing $Nrx1\beta$

Next we sought to determine if common clinical strains express SdrC and if the SdrC displayed by these isolates are capable of binding $Nrx1\beta$. Western blotting analysis of cell wall extractions from USA300, MW2 and MRSA252 display SdrC on their surface during exponential phase of growth in BHI (Fig. 5A1) and RPMI 10% FBS media, and when grown on TSB sheep blood

agar (data not shown). SdrC was not found in cell wall extractions from cells in stationary phase of growth (Fig. 5A2). We reasoned that SdrC absence from the cell wall extractions in the late stages of growth might be due to transcription cessation and proteolytical release from the cell wall. Western blotting analysis of concentrated culture supernatants revealed that SdrC was not detected in the supernatant during exponential growth but that all strains tested released SdrC fragments during late stationary growth phase (Fig. 5B1 and Fig. 5B2). SdrC fragments released in the supernatant migrated slower than recombinant SdrC_{N2N3} on the SDS-PAGE gels suggesting that a larger fragment of the protein is released from the cell wall. Immunoblotting analysis demonstrated that these fragments were recognized by antibodies raised against recombinant N2N3 domains and B-repeats (Fig. 5B3). USA300,




Figure 4. Determination of equilibrium constant of binding. (A) SdrC_{N2N3} binds specifically to β-neurexin 1. Increasing concentrations of SdrC_{N2N3} or ClfB_{N2N3} (black circle) were incubated with 10 nM fluorescein-labelled spNrx_{wt} peptide (black square) or spNrx_{scr} peptide (black triangle) for 3 hours at room temperature in the dark. Equation $\Delta P = \Delta P_{max} [protein] / (K_D + [protein])$ was used to calculate the equilibrium constant. The values from three experiments returned a $K_D = 2.50 \pm 0.49 \times 10^{-7}$ M. (B) SdrC_{N2N3} (0.5 μM) (black triangle) was incubated with increasing concentration of unlabelled spNrx_{wt} peptide (white circle) or spNrx_{scr} peptide (black circle) at room temperature. After 3 hours, the mixture was incubated for another 3 hours at room temperature in the dark with 10 nM fluorescein labeled spNrx_{wt} peptide. Fluorescence polarization was determined as described above. The values represented here are the mean ± SD of triplicates from three experiments.
doi:10.1371/journal.ppat.1000726.g004

MW2 and Newman display and release similar SdrC segments whereas MRSA252 expresses different SdrC species. One possible explanation for this difference may lie in the differences between their amino acid sequences (Fig. S5). These fragments were probed in ligand affinity western blots with recombinant Nrx₄₇₋₂₅₅ protein. Our data indicated that all strains tested release an active form of the SdrC protein capable to interact with Nrx1β (Fig. 5B4). A control membrane not probed with Nrx₄₇₋₂₅₅, did not indicate cross reactivity of anti-GST serum with SdrC fragments released into the supernatant (Fig. 5B5).

We next tested whether cell wall anchored SdrC mediates bacterial adherence to Nrx1β-expressing eukaryotic cells. We chose to express both proteins in heterologous systems that do not display any known interaction proteins. The full-length sdrC gene was cloned in pNZ8037 under a nisin inducible promoter. The plasmid was introduced in *L. lactis* for protein expression. Expression of SdrC on the bacterial cell surface was determined using immunofluorescence microscopy (data not shown). The full length Nrx1β gene was cloned in pCMV5 to obtain the translational fusion Nrx1β-mCherry [26] and a mutant FLAG-Nrx1β-mCherry where the SdrC binding site was replaced with a FLAG tag. These vectors were transfected into CHOK1 cells which were allowed to grow for 36 hours to ensure protein expression. Transiently transfected cells were incubated with *L. lactis* transformed with the vector alone, *L. lactis* expressing SdrC, *S. aureus* Newman, *S. aureus* Newman sdrC::Em' (DU 5988), USA300, MW2, or MRSA252 and bacterial attachment to the cultured mammalian cells were followed. Our results indicated that ten to twenty percent of the microorganisms expressing SdrC adhere to CHOK1 cells expressing wild-type Nrx1β. In contrast, we did not detect attachment when bacteria were incubated with untransfected cells or cells transiently transfected with the inactive FLAG-Nrx1β-mCherry or when Nrx1-mCherry expressing cells were incubated with the *S. aureus* mutant DU5988 or *L. lactis* transformed with an empty vector (Fig. 5C). Fluorescence micrographs revealed that bacteria colocalize with Nrx1β (Fig. S4).

Discussion

The success of *S. aureus* as a pathogen partly depends on the organism ability to effectively colonize different tissues in the host [9,27], evade host defense systems [28] and resist antibiotic therapy [29]. The molecular mechanisms involved in the development of different staphylococcal infections are incompletely understood but likely involve a large set of virulence factors. Many of these virulence factors interact with and manipulate specific molecular targets in the host. To interact with host targets most virulence factors are either secreted or exposed on the surface of the bacteria. Gram-positive bacteria express a family of surface proteins that are covalently anchored to the cell wall at their C-terminus. The amino acid sequences of these CWA proteins vary dramatically from one organism to another. Nevertheless, structural predictions identify a subgroup of CWA proteins with a common structural organization. This subgroup includes the known staphylococcal MSCRAMMs but also several proteins with unknown function. We hypothesize that the latter also act as adhesins and interact with molecules in the host [7,9]. In our search for ligands for these orphan adhesins, we here report that screening a phage display library of linear peptides is a useful strategy to identify host targets for orphan MSCRAMMs. To demonstrate the validity of this approach we first used SdrG_{N2N3} to pan a commercially available library of 12 amino acid inserts in the P3 phage protein. The consensus sequence identified from this analysis overlapped with the previously determined SdrG binding site located in the N-terminus of the Fg β chain. We subsequently predicted a N2N3 domain in SdrC and used a recombinant form of this segment as a target in panning the peptide library. A pattern search using the degenerate consensus sequence returned by the phage display experiment against the whole human proteome identified β-neurexin isoforms as the putative binding partners for SdrC. The human proteome contains three β-neurexin proteins, which share approximately 60% identity with each other [14]. Each β-neurexin isoform contains a variant of the peptide sequence identified by phage display located close to the amino-terminal end of the protein, which is exposed on the surface




Figure 5. SdrC mediates *S. aureus* clinical isolates adherence to Nr1β expressing cells. (A) Western immunoblotting to detect SdrC expression on the surface of different bacterial strains. *S. aureus* Newman, DU5988 and clinical strains USA300, MW2, MRSA252 were grown to exponential phase (A1) or stationary phase (A2) in BHI. CWA proteins extracted as described in material and methods were separated on 4–15% gradient SDS-PAGE gradient gels, transferred to nitrocellulose and probed with anti-SdrC serum. (B) Culture supernatants from the above mentioned strains grown to exponential (B1) or stationary phase (B2) were concentrated and separated on 4–15% gradient SDS-PAGE gradient gels after protein A removal. After gel separation the proteins were transferred to nitrocellulose and probed with anti-SdrC serum (B1 and B2). The nitrocellulose blot corresponding to the stationary phase supernatants was stripped and re-probed with anti-SdrC B-repeats antibody (B3). The same blot was stripped again and probed with NrX₄₇₋₂₅₅. The binding of NrX1β was detected with anti-GST serum (B4). A second membrane (B5) was not incubated with NrX₄₇₋₂₅₅ and therefore served as control for the primary and secondary sera. (C) *L. lactis* empty vector, *L. lactis* SdrC, *S. aureus* Newman, DU5988 (*S. aureus* Newman *sdrC::Em*), USA300, MW2 or MRSA252 were incubated with CHOK1 transiently transfected with NrX1β-mCherry or FLAG NrX1β-mCherry grown in 24 well plates. Attachment of bacteria was reported as a percentage of total bacteria at the end of incubation. Values shown represent mean ± SD of three experiments (*p ≤ 0.001). doi:10.1371/journal.ppat.1000726.g005

of the eukaryotic cell and therefore is accessible for binding. Moreover, this sequence starts nine residues after the first amino acid of β-neurexin, which is particularly interesting because the SdrC_{N2N3} binding site in Fg also maps nine residues from the N-terminus of the β chain polypeptide.

Next, we demonstrated that both the SdrC_{N2N3} and the SdrC_A fragments bind with high affinity to the amino-terminal segment of NrX1β, only when the polypeptide contains the amino acid sequence identified by phage display. The interaction does not promote bacterial internalization (Fig. S1), the K_d of interaction is not influenced by the amount of protein coated on the plates (Fig. S2) and it is not influenced by metal ions (Fig. S3). We further

demonstrated that a synthetic peptide corresponding to the binding site in β-neurexin effectively inhibits the binding of SdrC to NrX1β. These data indicated that recombinant SdrC_{N2N3} is able to bind recombinant NrX1β and that the binding site is the amino acid sequence identified by phage display. In this study, we have also shown that the binding of SdrC to NrX_{wt} peptide exhibits a K_D of 2.5 × 10⁻⁷, which is comparable to those recorded for high affinity CWA protein/host protein interactions. Further, we showed that *S. aureus* Newman, USA300, MW2 and MRSA252 release a fragment of SdrC capable of binding NrX1β. Heterologous expression of SdrC on the surface of *L. lactis* mediates bacterial attachment to NrX1β transiently expressed on CHOK1

cells. Similarly, clinical strains expressing SdrC adhere to wild-type Nr1β heterologously expressed in CHOK1. This experiment demonstrates that SdrC can interact with Nr1β when the two proteins are appropriately expressed on bacteria and mammalian cells, respectively.

The biological significance of the SdrC-Nr1β interaction is unclear. mRNA for β-neurexins is found in many tissues [30] but the protein was only detected in neuronal tissues. It is possible that Nr1β protein expression occurs in non-neuronal cells or tissues that have not yet been examined and/or that protein expression in non-neuronal tissues is induced under certain conditions. SdrC may serve as an adhesin mediating bacterial attachment to Nr1β in staphylococcal infections of neuronal tissues or where the target protein is expressed. It is also possible that during the course of the infection biologically active fragments could be released from the bacteria during proteolytic processing of CWA proteins. SdrC_{N2N3} fragments which are released and resistant to further proteolytic degradation could interact with Nr1β at tissues distant from the primary infection site. In this scenario it would be interesting to examine if SdrC fragments can induce intracellular signaling through its interaction with Nr1β or interfere with the neurexin-neurologin interaction.

A review of the literature reveals that staphylococcal endocarditis and sepsis have been associated with polyneuropathy and in several rare cases even with reversible acute tetraplegia. Studies suggest that polyneuropathy may often follow after sepsis [31,32,33]. Before recent advances in medicine, such as life support in the ICUs, septic death likely occurred before the neuromuscular signs could be observed. The most common manifestations of polyneuropathy are difficulty in weaning from the ventilator and limb weakness. Electrophysiologic and histopathologic investigations demonstrated axonal degeneration of motor and sensory fibers that is different from autoimmune syndromes that can develop during infection. The etiology of sepsis-associated polyneuropathy is unclear. It was thought to be caused by neuromuscular blocking agents used to ease mechanical ventilation, antibiotic toxicity, corticosteroids, microanoxia, nutritional deficiency, and high levels of inflammatory mediators. However, the cause of the disease remains unclear and is likely due to a combination of the above-mentioned factors (reviewed in [31]). Moreover, the molecular mechanism of axonal degeneration in sepsis-associated polyneuropathy is unknown. On the other hand, studies investigating neurotransmission, revealed that inhibition of Nr1β-Nlg interaction led to late-onset neuromuscular deterioration [34]. These observations raise the possibility that SdrC interaction with Nr1β may contribute to sepsis-associated polyneuropathy. Future studies will investigate in detail the ability of SdrC to interfere with neurexin biology *in vitro* and *in vivo*.

Materials and Methods

Media and growth conditions

E. coli was cultured in LB medium (Sigma, St. Louis, MO) containing ampicillin (100 µg/ml) at 37°C with shaking at 250 rpm. *S. aureus* was cultured in tryptic soy broth at 37°C with shaking at 250 rpm. *L. lactis* was cultured in GM17 (Oxoid, LTD, Hampshire, UK) containing 0.5% glucose and chloramphenicol (10 µg/ml) (Sigma, St. Louis, MO) at 30°C without shaking. To express SdrC in *L. lactis*, overnight cultures were diluted 1:100, grown for another 3 hours and induced with nisin (1.6 ng/ml) overnight, unless otherwise mentioned. CHOK1 cells were grown in CD CHO media (GIBCO, Grand Island, NY) at 37°C in a humidified chamber with 5% CO₂.

Bacterial strains used for adhesion

S. aureus strains MW2, USA300 and MRSA were from NARSA or previously described [35]. The *sdrC* null mutant (*sdrC*:pG⁺Host) of *S. aureus* Newman (DU5988) was described previously [36]. The *sdrC* gene was cloned into the nisin-inducible expression vector pNZ8037 [37] and transferred into *L. lactis* NZ9800 by previously described procedures [38].

Molecular modeling

Modeling of SdrC was performed using PHYRE Protein Homology/analogY Recognition Engine Version 2.0 (<http://www.sbg.bio.ic.ac.uk/phyre/html/index.html>)

Plasmid construction

DNA manipulation was performed using standard methods. DNA modification and restriction enzymes were purchased from New England Biolabs, Inc. (Ipswich, MA) or Promega (Madison, WI) and used according to the manufacturer protocol. Fragments encoding different domains of SdrC or Nr1β were amplified by PCR from *S. aureus* genomic DNA, plasmids pCMV5 Nr1β-Fc or pCMV5 FLAG-Nr1β-mCherry [26] and oligonucleotide primers listed in Table 4. The PCR products were analyzed by agarose gel electrophoresis and purified using a QIAquick gel extraction kit (Qiagen, Sciences, MD). To construct SdrC expression plasmids, a BamHI-HindIII fragment containing the appropriate gene segment was cloned into pQE30 (Qiagen, Sciences, MD). To construct the Nr1β-mCherry translational fusion, we first amplified the N-terminal 765 nt of Nr1β from pCMV5 Nr1β-Fc to obtain a PCR product corresponding to the first 255 aa of neurexin protein (including the signal sequence). Second, we amplified the Nr1β 255-mCherry from plasmid pCMV5 FLAG-Nr1β-mCherry to obtain the translational fusion of the 3' end fragment of neurexin gene with mCherry from *Discosoma* sp. The newly synthesized PCR fragments were used as a template for an overlapping PCR to obtain full length neurexin 1β-mCherry as a translational fusion. A BglII-XbaI fragment containing the overlapping PCR was cloned into pCMV5.

To obtain GST-tagged expression proteins we amplified Nr1₄₇₋₂₅₅ (aa) or Nr1₇₀₋₂₅₅ (aa) from pCMV5 Nr1β-Fc using primers described in Table 1. After purification we cloned these

Table 4. Oligonucleotide primers used to amplify gene fragments.

Expression construct	Primer sequence
SdrC ₅₂₋₄₉₆ (SdrC _A)	5' CGCAGGATCCGAGAACATACGAATGGAG
	5' CGCAAAGCTTACTTTTGGTCGCCATTAGCAC
SdrC ₁₇₈₋₄₉₆ (SdrC _{N2N3})	5' CCCGGATCCGGAACAATGTTAATGATAAAGTACAT
	5' CCCAAGCTTTTATTCTTTTGGTCGCCATTAG
Nr1 ₄₇₋₂₅₅	5' CACCATGGCATCCAGTTGGGAGCCG
	5' TTTACATTTCCCACTATGGCGATG
Nr1 ₇₀₋₂₅₅	5' CACCATGAGTTGGGAGCCGACCACA
	5' TTTACATTTCCCACTATGGCGATG
Nr1 ₁₋₂₅₅	5' CCGCAGATCTATGTACAGAGGATGCTCCGGT
	5' TTTACATTTCCCACTATGGCGATG
Nr1 _{255-mCherry}	5' CATCGCCATAGTGGGAAATGTGAT
	5' GCGCTCTAGATTACTTGTACAGCTCGTCCATGCCG

The enzyme restriction sites are underlined.
doi:10.1371/journal.ppat.1000726.t004

fragments in the *E. coli* expression system with Gateway Technology (Invitrogen, Inc, Carlsbad, CA) according to the manufacturer's instructions. All plasmid constructs were sequenced to ensure the integrity of the amplified fragments (Baylor College of Medicine DNA Sequencing Core Facility).

Protein expression and purification

Plasmids pQE30-SdrC₅₂₋₄₉₆ and pQE30-SdrC₁₇₈₋₄₉₆ were transformed into *E. coli* Topp 3. Overnight starter cultures were diluted 1:50 in LB containing ampicillin (100 μ g/ml) and incubated with shaking until the culture reached OD₆₀₀ 0.6–0.8. Protein expression was induced by adding 0.1 mM IPTG (final concentration) and continuing the incubation for 4 hours. Bacterial cells were harvested by centrifugation, resuspended in PBS and frozen at -80°C . Plasmids pDEST Nrx₄₇₋₂₅₅ and pDEST Nrx₇₀₋₂₅₅ were transformed into BL21-AI (Invitrogen, Inc, Carlsbad, CA). Cultures were induced with 0.1% arabinose for 16 h at room temperature. Bacterial cells were harvested by centrifugation, resuspended in PBS containing EDTA-free Complete Protease Inhibitor (Roche Diagnostics, Mannheim, Germany) and frozen at -80°C .

To purify SdrC₅₂₋₄₉₆ and SdrC₁₇₈₋₄₉₆, cells containing recombinant protein fragments were passed through a French press (1100 p.s.i.). Cellular debris was removed by centrifugation at 40,000 rpm for 20 minutes followed by filtration through a 0.45 μ M membrane. The filtered bacterial lysate were applied at 2 ml/min on a 5 ml nickel-charged HiTrap Chelating column (GE Healthcare, Uppsala, Sweden) equilibrated with 10 mM Tris HCl, 100 mM NaCl pH 7.9. The column was washed with 40 volumes of 10 mM Tris HCl, 100 mM NaCl, 20 mM imidazole. Bound protein was eluted with a linear gradient of imidazole (10 to 200 mM, total volume 200 ml) yielding proteins that were >95% pure.

Fractions containing the SdrC 52–496 recombinant protein were dialyzed overnight in 4 L of 25 mM Tris-Cl pH 7.9 containing 10 mM EDTA and 1 mM 1,10 O-phenantroline. The dialysed sample was applied at 2 ml/minute to a MonoQ anion exchange column equilibrated with 25 mM Tris-Cl pH 7.9 for further purification. Recombinant protein of interest was collected from the flow through and dialyzed against PBS pH 7.4. Sample was reappplied on a nickel-charged HiTrap Chelating column. Pure protein was dialyzed against 50 mM EDTA to remove excess nickel and then in HBS (10 mM HEPES pH 7.4, 150 mM NaCl, 3 mM EDTA) to remove excess EDTA.

Fractions containing the SdrC₁₇₈₋₄₉₉ recombinant protein were dialyzed overnight in 25 mM Tris-Cl pH 8.7. Dialyzed sample was applied at 2 ml/minute to a MonoQ anion exchange column equilibrated with 25 mM Tris-Cl pH 8.7. Bound protein was eluted with a linear gradient of NaCl (0–1 M NaCl, 160 ml). Fractions containing pure protein were dialyzed against 50 mM EDTA to remove excess nickel and then against HBS to remove excess EDTA.

To purify neurexin GST-tagged fragments, cells were passed through a French press (1100 p.s.i.) in the presence of Complete protease inhibitor tablets (Roche Molecular Biochemicals). Cellular debris was removed as described above. Clear cell lysate was applied to a glutathione-Sepharose column (Sigma, St. Louis, MO). Bound protein was eluted with 10 nM glutathione pH 8. Recombinant protein was concentrated and applied to a S-75 Sephadex (GE Healthcare, Uppsala, Sweden). Proteins were eluted with PBS pH 7.4.

Mass spectrometry analysis and N-terminal sequencing were performed at Tufts Protein Core Facility, Tufts University.

SDS-PAGE and Western

Recombinant proteins were analyzed by SDS-PAGE using standard procedure [39] on 12% acrylamide gels. Gels were stained with Coomassie Blue. CWA proteins from cells grown to exponential (OD₆₀₀ = 8) or stationary phase (10 h) were harvested, washed 3 times with PBS and concentrated to OD₆₀₀ = 50. CWA proteins were solubilized with 200 μ g/ml lysostaphin in a buffer designed to minimize bacterial lysis composed of 30% sucrose, 50 mM Tris/Base pH 8, 20 mM MgCl₂ and 50 μ g/ml protease inhibitor (Roche) for 30 min at 37°C. Cell wall fractions were then incubated with 100 μ l Sepharore-IgG for 3 h at 4°C to remove protein A. Culture supernatants were concentrated to OD₆₀₀ = 10 after the addition of 100 μ g/ml protease inhibitor (Roche). Protein A was removed by incubating with 100 μ l Sepharore-IgG. Extracts or supernatants were separated on 4–15% SDS-PAGE gels. For Western blotting, proteins were transferred electrophoretically to nitrocellulose or polyvinylidene difluoride membranes (Bio-Rad, Hercules, CA) using the semi-dry system (Bio-Rad, Hercules, CA) in Tris (0.02 M)-Glycine (0.15 M)-methanol (20%) buffer. Membranes were blocked in 10% milk and incubated with the appropriate serum. Briefly, membranes A1, A2, B1 and B2 were incubated with rabbit anti-SdrC_{N2N3} serum (H.T.I. BioProducts, Inc) and the appropriate secondary serum, which is listed below. B2 membrane was stripped (Restore western blot stripping Buffer, Pierce, Rockford, IL) for 10 minutes at room temperature and then re-probed with a rabbit anti-SdrC B-repeats serum (H.T.I. BioProducts, Inc) (B3). The membrane was stripped a second time and incubated for 1 hour at room temperature with Nrx₄₇₋₂₅₅ recombinant protein, followed by goat anti-GST serum. (Invitrogen, Carlsbad, CA) (B4). A control membrane identical with B2 was probed with the same sera but was not incubated with Nrx₄₇₋₂₅₅ recombinant protein (B5). HRP-labeled goat antibody (1:5000) or HRP-labeled goat anti-mouse or HRP-labeled rabbit anti-goat (BioRad, Hercules, CA) were used to detect bound primary antibody by incubating for 1 hour at room temperature. Membranes were developed with ECL reagent (Pierce, Rockford, IL), exposed to Kodak X-ray film.

Phage display

NEB Ph.D.12-mer random library was incubated with immobilized SdrC N2N3, SdrG N2N3 or BSA at 1 μ g/well. The input was 1.5×10^{11} transducing units with a complexity of 2.7×10^9 electroporated sequences. The assay was performed according to the manufacturer's instructions. The output phage plaques were 10^4 transducing units. Wells were washed with 0.5% Tween in TBS buffer, blocked with 10% BSA and incubated with 1×10^8 transducing units of each amplified phage or insertless phage (fd-tet). To remove unbound phage, wells were washed 10 times with 0.5% Tween in TBS. Bound phage was detected with anti-M13-HRP antibody (GE Healthcare, Buckinghamshire, UK).

Solid phase binding assay

Each well of Immulon 4BH plates was coated with 1 μ g of recombinant Nrx₄₇₋₂₅₅-GST, Nrx₇₀₋₂₅₅-GST or GST overnight at 4°C. Coated wells were blocked for 1 hour at room temperature with 2% BSA in 0.05% Tween-TBS buffer. Increasing concentrations of SdrC_{N2N3} or A-region were added to the wells and incubated for 1 hour at room temperature. Bound protein was detected with a polyclonal antibody (1:3000) against SdrC_{N2N3} (H.T.I. BioProducts) followed by an anti-rabbit HRP-labeled antibody (1:5000) (Bio-Rad Labs, Hercules, CA). Color development was performed using SigmaFast OPD (Sigma, St. Louis, MO) and the binding was measured using a microtiter plate reader (Molecular Devices) at 450 nm. For inhibition assays, 0.5 μ M

SdrC₁₇₈₋₄₉₆ protein was incubated with increasing concentrations of peptides prior to the solid phase binding assay. Data presented represent the mean±SD of three independent experiments performed in triplicate. The binding was analyzed by non-linear regression for one binding site (GraphPad Prism).

Fluorescence polarization

Nrx_{wt} and Nrx_{scr} peptides were labeled with fluorescein as described previously [10]. Increasing concentrations of SdrC₁₇₈₋₄₉₆ were incubated with 10 nM labeled peptide for 3 hours in the dark at room temperature. Polarization was measured using Luminescence Spectrometer LB50B (Perkin Elmer). The data were analyzed by non-linear regression for one binding site. The equilibrium dissociation constant was calculated using the equation $\Delta P = \Delta P_{max} \times [protein] / (K_D + [protein])$ (Eq.1) where ΔP is the change in fluorescence polarization, ΔP_{max} is the maximum change in fluorescence polarization and K_D is the equilibrium dissociation constant of the interaction.

For inhibition assays, 0.5 μM SdrC₁₇₈₋₄₉₆ was first incubated with increasing concentrations of unlabeled Nrx_{wt} or unlabeled Nrx_{scr} for 3 hours at room temperature. The mixture was then incubated for 3 hours with fluorescein-labeled Nrx_{wt}. The results presented represent the mean±SD of three independent experiments.

Adherence assay

CHOK1 cells were grown in 24-well tissue culture plates to approximately 80% confluence. Cells were transfected with either pCMV5 Nrx1β-mCherry or pCMV5 FLAG-Nrx1β-mCherry using Lipofectamine 2000 (Invitrogen, Carlsbad, CA) according to the manufacturer's instructions and grown for 36 hours to allow for protein expression. For attachment assays, bacteria were washed 2 times in PBS, resuspended in CD CHO media and incubated for 1 h with CHOK1 cells at an MOI of 10. To remove unattached bacteria, cells were washed 4 times with PBS. To determine the number of attached bacteria, CHOK1 cells were lysed with sterile water and serial dilutions plated on the appropriate media. The number of attached bacteria was reported as a percent of total bacteria at the end of the incubation period. Each experiment was performed in triplicate wells and repeated three times. Statistical analysis was performed using the Student's t-test.

Protein accession numbers (UniProt) SdrC

S. aureus Newman - O86487; *S. aureus* USA300 - A8YZQ9; *S. aureus* MW2 - Q8NXX7; *S. aureus* MRSA252 - Q6GJA7. Neurexin1β: P58400.

References

- Lowy FD (1998) Staphylococcus aureus infections. N Engl J Med 339: 520–532.
- Clarke SR, Foster SJ (2006) Surface adhesins of *Staphylococcus aureus*. Adv Microbiol Physiol 51: 187–224.
- Patti JM, Allen BL, McGavin MJ, Hook M (1994) MSCRAMM-mediated adherence of microorganisms to host tissues. Annu Rev Microbiol 48: 585–617.
- Patti JM, Hook M (1994) Microbial adhesins recognizing extracellular matrix macromolecules. Curr Opin Cell Biol 6: 752–758.
- Marraffini LA, Dedent AC, Schneewind O (2006) Sortases and the art of anchoring proteins to the envelopes of gram-positive bacteria. Microbiol Mol Biol Rev 70: 192–221.
- Ton-That H, Marraffini LA, Schneewind O (2004) Protein sorting to the cell wall envelope of Gram-positive bacteria. Biochim Biophys Acta 1694: 269–278.
- Ponnuraj K, Bowden MG, Davis S, Gurusiddappa S, Moore D, et al. (2003) A “dock, lock, and latch” structural model for a staphylococcal adhesin binding to fibrinogen. Cell 115: 217–228.
- Deivanayagam CC, Wann ER, Chen W, Carson M, Rajashankar KR, et al. (2002) A novel variant of the immunoglobulin fold in surface adhesins of *Staphylococcus aureus*: crystal structure of the fibrinogen-binding MSCRAMM, clumping factor A. EMBO J 21: 6660–6672.

Supporting Information

Table S1 Pattern search results

Found at: doi:10.1371/journal.ppat.1000726.s001 (0.06 MB PDF)

Figure S1 Thirty-six hours prior to infection, CHOK1 cells where transfected with Nrx1β-mCherry. Bacteria were added to each well at MOI of 10 and incubated for 3 hours in a humidified incubation chamber at 37°C with 5% CO₂. To remove unbound bacteria, wells were washed 3 times with PBS. After washing, cells were incubated in DMEM containing gentamicin (100 μg/ml) for 1 h. Cells were washed 3 times in PBS, lysed with water and dilution plated. Internalized bacteria where represented as percent of total bacteria at the end of the incubation period.

Found at: doi:10.1371/journal.ppat.1000726.s002 (0.10 MB PDF)

Figure S2 The solid phase binding assay was performed as described in Materials and Methods. The wells were coated with 0.5 μg/well Nrx₄₇₋₂₅₅.

Found at: doi:10.1371/journal.ppat.1000726.s003 (0.10 MB PDF)

Figure S3 The solid phase binding assay was performed as described in Materials and Methods. The wells were coated with 1 μg/well Nrx₄₇₋₂₅₅. EDTA, [Ca²⁺], [Mg²⁺], [Mn²⁺] were added to 1 μM SdrC_{N2N3} prior incubation with Nrx1β at a final concentration of 10 μM.

Found at: doi:10.1371/journal.ppat.1000726.s004 (0.10 MB PDF)

Figure S4 CHOK1 cells transiently transfected with Nrx1β-mCherry or FLAG Nrx1β-mCherry and grown on glass coverslips were incubated with either *L. lactis* empty vector, *L. lactis* SdrC, *S. aureus* Newman or DU5988 (*S. aureus* Newman *sdrC::Emr*) Nrx1β expression was monitored by the red fluorescence due to mCherry. Bacterial cells were detected with anti-SdrC antibodies and FITC-labeled secondary antibodies (green). CHOK1 nuclei were stained with DAPI (blue). Attachment of bacteria to Nrx1β-expressing CHOK1 cells is shown (merged).

Found at: doi:10.1371/journal.ppat.1000726.s005 (2.28 MB PDF)

Author Contributions

Conceived and designed the experiments: EMB VKG MH. Performed the experiments: EMB. Analyzed the data: EMB VKG RCM MH. Contributed reagents/materials/analysis tools: SG TJF TCS. Wrote the paper: EMB RCM MH.

15. Ushkaryov YA, Petrenko AG, Geppert M, Sudhof TC (1992) Neurexins: synaptic cell surface proteins related to the alpha-latrotoxin receptor and laminin. *Science* 257: 50–56.
16. Mullen LM, Nair SP, Ward JM, Rycroft AN, Henderson B (2006) Phage display in the study of infectious diseases. *Trends Microbiol* 14: 141–147.
17. Sergeeva A, Kolonin MG, Molldrem JJ, Pasqualini R, Arap W (2006) Display technologies: application for the discovery of drug and gene delivery agents. *Adv Drug Deliv Rev* 58: 1622–1654.
18. Kolonin MG, Bover L, Sun J, Zurita AJ, Do KA, et al. (2006) Ligand-directed surface profiling of human cancer cells with combinatorial peptide libraries. *Cancer Res* 66: 34–40.
19. Smith GP, Petrenko VA (1997) Phage Display. *Chem Rev* 97: 391–410.
20. Deivanayagam CC, Perkins S, Danthuluri S, Owens RT, Bice T, et al. (1999) Crystallization of ClfA and ClfB fragments: the fibrinogen-binding surface proteins of *Staphylococcus aureus*. *Acta Crystallogr D Biol Crystallogr* 55: 554–556.
21. Ganesh VK, Rivera JJ, Smeds E, Ko YP, Bowden MG, et al. (2008) A structural model of the *Staphylococcus aureus* ClfA-fibrinogen interaction opens new avenues for the design of anti-staphylococcal therapeutics. *PLoS Pathog* 4: e1000226. doi:10.1371/journal.ppat.1000226.
22. Ni Eidhin D, Perkins S, Francois P, Vaudaux P, Hook M, et al. (1998) Clumping factor B (ClfB), a new surface-located fibrinogen-binding adhesin of *Staphylococcus aureus*. *Mol Microbiol* 30: 245–257.
23. Hartford OM, Wann ER, Hook M, Foster TJ (2001) Identification of residues in the *Staphylococcus aureus* fibrinogen-binding MSCRAMM clumping factor A (ClfA) that are important for ligand binding. *J Biol Chem* 276: 2466–2473.
24. McAleese FM, Walsh EJ, Sieprawska M, Potempa J, Foster TJ (2001) Loss of clumping factor B fibrinogen binding activity by *Staphylococcus aureus* involves cessation of transcription, shedding and cleavage by metalloprotease. *J Biol Chem* 276: 29969–29978.
25. Ushkaryov YA, Hata Y, Ichtchenko K, Moomaw C, Afendis S, et al. (1994) Conserved domain structure of beta-neurexins. Unusual cleaved signal sequences in receptor-like neuronal cell-surface proteins. *J Biol Chem* 269: 11987–11992.
26. Boucard AA, Chubykin AA, Comoletti D, Taylor P, Sudhof TC (2005) A splice code for trans-synaptic cell adhesion mediated by binding of neuroligin 1 to alpha- and beta-neurexins. *Neuron* 48: 229–236.
27. Bubeck Wardenburg J, Patel RJ, Schneewind O (2007) Surface proteins and exotoxins are required for the pathogenesis of *Staphylococcus aureus* pneumonia. *Infect Immun* 75: 1040–1044.
28. Foster TJ (2005) Immune evasion by staphylococci. *Nat Rev Microbiol* 3: 948–958.
29. Chambers HF (2005) Community-associated MRSA—resistance and virulence converge. *N Engl J Med* 352: 1485–1487.
30. Occhi G, Rampazzo A, Boffagna G, Antonio Danieli G (2002) Identification and characterization of heart-specific splicing of human neurexin 3 mRNA (NRXN3). *Biochem Biophys Res Commun* 298: 151–155.
31. Chen HC, Tsai CS, Lee JT, Chen CA, Chang FY (2005) Acute quadriplegia complicating critical illness polyneuropathy in a patient with infective endocarditis: a case report. *J Infect* 50: 153–157.
32. Roder BL, Wandall DA, Espersen F, Frimodt-Moller N, Skinhoj P, et al. (1997) Neurologic manifestations in *Staphylococcus aureus* endocarditis: a review of 260 bacteremic cases in nondrug addicts. *Am J Med* 102: 379–386.
33. Caksen H, Uner A, Arslan S, Anlar O, Odabas D, et al. (2004) Severe peripheral polyneuropathy in a child with infective endocarditis caused by *Staphylococcus aureus*. *Acta Neurol Belg* 104: 114–116.
34. Andres C, Beerli R, Friedman A, Lev-Lehman E, Henis S, et al. (1997) Acetylcholinesterase-transgenic mice display embryonic modulations in spinal cord choline acetyltransferase and neurexin Ibeta gene expression followed by late-onset neuromotor deterioration. *Proc Natl Acad Sci U S A* 94: 8173–8178.
35. Voyich JM, Otto M, Mathema B, Braughton KR, Whitney AR, et al. (2006) Is Panton-Valentine leukocidin the major virulence determinant in community-associated methicillin-resistant *Staphylococcus aureus* disease? *J Infect Dis* 194(12): 1761–70.
36. O'Brien L, Kerrigan SW, Kaw G, Hogan M, Penades J, et al. (2002) Multiple mechanisms for the activation of human platelet aggregation by *Staphylococcus aureus*: roles for the clumping factors ClfA and ClfB, the serine-aspartate repeat protein SdrE and protein A. *Mol Microbiol* 44: 1033–1044.
37. de Ruyter PG, Kuipers OP, de Vos WM (1996) Controlled gene expression systems for *Lactococcus lactis* with the food-grade inducer nisin. *Appl Environ Microbiol* 62: 3662–3667.
38. Loughman A, Fitzgerald JR, Brennan MP, Higgins J, Downer R, et al. (2005) Roles for fibrinogen, immunoglobulin and complement in platelet activation promoted by *Staphylococcus aureus* clumping factor A. *Mol Microbiol* 57: 804–818.
39. Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227: 680–685.