

## BRIEF COMMUNICATION

# Anorexia, bulimia, and obesity: Shared decision making deficits on the Iowa Gambling Task (IGT)

AMY BROGAN,<sup>1</sup> DAVID HEVEY,<sup>1</sup> AND RICCARDO PIGNATTI<sup>2</sup>

<sup>1</sup>School of Psychology, Trinity College Dublin, Dublin 2, Ireland

<sup>2</sup>Istituto Auxologico Italiano, Neuropsychology Unit, Piancavallo (VB), Italy

(RECEIVED October 22, 2009; FINAL REVISION March 8, 2010; ACCEPTED March 9, 2010)

### Abstract

The pathological eating behaviors in Anorexia Nervosa (AN), Bulimia Nervosa (BN), and obesity are characterized by a preference for high immediate reward, despite higher future losses in terms of both physical and psychological outcomes. The present study compared the decision making profile of females with a diagnosis of AN ( $n = 22$ ), BN ( $n = 17$ ), obesity ( $n = 18$ ), and a healthy weight comparison group ( $n = 20$ ) using a standardized neuropsychological test, the Iowa Gambling Task (IGT). The three clinical groups (AN, BN, obesity) were significantly impaired on the IGT compared with the comparison group on both overall task performance and task learning; however, the three clinical groups were not significantly different from each other. Sixty-one percent to 77% of the clinical groups reached the threshold for impairment on the IGT, compared with 15% of the comparison group. The potential basis for this shared decision making profile is discussed. (*JINS*, 2010, *16*, 711–715.)

**Keywords:** Neuropsychology, Prefrontal cortex, Cognition, Choice behavior, Reward, Eating disorders

## INTRODUCTION

### Pathological Eating Behavior – Immediate Versus Long-Term Reward

Anorexia Nervosa (AN) and Bulimia Nervosa (BN) are characterized by the tendency to make decisions that have positive short-term consequences, but that may result in long-term negative outcomes (Brand, Franke-Sievert, Jacoby, Markowitsch, & Tuschen-Caffier, 2007). In AN, individuals severely restrict their caloric intake to obtain an immediate reward (the relief of anxiety elicited by food phobia) while ignoring the progressive and inevitable decline in their physical health (Cavedini et al., 2004). Similarly, in BN, there is a tendency to engage in self-damaging behaviors (binge eating/purging) that provide immediate benefit (relief of tension), at the cost of negative medical consequences (Boeka & Lokkenz, 2006). This diminished ability to assess future consequences has also been hypothesized as characteristic of overweight

and obese individuals (Davis, Levitan, Mugllia, Bewell, & Kennedy, 2004). These individuals choose to overeat (often palatable energy-dense foods) despite the long-term negative health consequences. Common to AN, BN, and obesity is an inability to appropriately regulate eating behavior.

### The Iowa Gambling Task (IGT) and Eating Pathology

The IGT (Bechara, Damasio, Damasio, & Anderson, 1994) was developed for functional assessment in ventromedial prefrontal cortex (VmpPFC) patients, who show impaired decision making. Card deck selection assesses the ability to forgo immediate gain for a longer-term successful outcome. It simulates decision making strategy by factoring uncertainty, reward, and punishment (Bechara et al., 1994). The IGT is underpinned by the Somatic Marker Hypothesis (SMH) (Damasio, 1994), a theory that views decision making as a dual combination of “high reason,” carrying out a logical cost-benefit analysis of a given action, and “somatic marker” biasing signals, indicating how rewarding or punishing an action is likely to be. Successful IGT performance has been correlated

Correspondence and reprint requests to: Amy Brogan, School of Psychology, Trinity College Dublin, Dublin 2, Ireland. E-mail: broganam@tcd.ie

with the development of “somatic marker” signals (as indexed by anticipatory skin conductance responses [SCRs]) in healthy control participants, and with the absence of such signals in VmPFC patients (Bechara, Tranel, Damasio, & Damasio, 1996). The IGT has been validated with a wide range of clinical groups (see Dunn, Dalgleish, & Lawrence, 2006).

Despite calls for direct comparisons of the decision making profile in disordered eating groups (Brand et al., 2007), research has assessed each group separately using the IGT. Significant impairment on the IGT was revealed in the acute stage of AN, independent of illness severity or Body Mass Index (BMI) (Cavedini et al., 2004). Female AN patients who had a better IGT decision making profile at baseline showed a significantly greater improvement in nutritional status after a cognitive-behavioral treatment program (Cavedini et al., 2006). Consistent with the SMH (Damasio, 1994), impairment in decision making ability in females with AN was associated with significantly attenuated somatic marker signals (indexed by SCRs), compared with both healthy controls and females with long-term recovery from AN (healthy BMI and minimum 1 year’s menstruation) (Tchanturia, Liao, Uher, Lawrence, Treasure, & Campbell, 2007). Significant impairment was found in undergraduate women with BN, wherein IGT performance negatively correlated with bulimic symptomatology (Boeka & Lokkenz, 2006). A recent study found that female BN patients performed poorly on the IGT (Liao et al., 2009), but with no impairment in SCR (in contrast to results with AN; Tchanturia et al., 2007). In a comparison of AN females (who were either underweight or recovered to normal weight) and BN females, research found no significant differences on the IGT between these eating disorder groups and controls (Bosanac et al., 2007). However, the eating disorder groups tended to score lower, and failure to reach significance may be attributable to low statistical power. Finally, both healthy women with a high BMI (BMI > 25) (Davis et al., 2004), and individuals with severe obesity (BMI > 34) (Pignatti, Bertella, Albani, Mauro, Molinari, & Semenza, 2006) performed disadvantageously on the IGT. A comparison of obese women with and without Binge Eating Disorder (BED) found that both groups had lower mean IGT scores than a normal weight group, but were not significantly different from each other. When education was included in the analyses, those with a higher education performed better and the group differences were not significant (Davis, Patte, Curtis, & Reid, 2010).

## Aims

Separate examinations of AN, BN, and obesity have revealed significant decision making impairments in terms of IGT performance. This study furthers this research by directly comparing the decision making profiles of AN, BN, and obese females on the IGT, and evaluating their performance in relation to a healthy comparison group.

## METHOD

### Participants and Procedure

#### *AN and BN participants*

Twenty-two female participants who met the DSM-IV-TR (American Psychiatric Association [APA], 2000) criteria for AN and 17 female participants who met the BN criteria were recruited from a rehabilitation program in an Italian hospital specialized in the treatment of obesity and eating disorders. The AN group ranged in age from 18 to 46 years, and the BN group from 19 to 38 years. All patients underwent diagnostic interviews conducted by a trained psychiatrist and a psychologist. Participants were not tested during an acute stage of illness.

#### *Obese participants*

Eighteen female obese individuals were recruited from an Irish hospital specialized in obesity management. Participants ranged in age from 30 to 73 years.

Exclusion criteria common to all three clinical groups were underage (<18 years), and failure to complete the compulsory education period (minimum 8 years of formal education). Additional exclusion criteria for the AN and BN groups were disease duration of <1 year, a comorbid personality disorder or other pervasive psychopathology, and oppositional behaviors during the experimental session (two participants removed). For the obese group, other exclusion criteria included a diagnosis of psychopathology, BED, or a documented history of binge eating by the hospital’s clinical psychologist.

#### *Comparison participants*

The comparison group comprised 20 normal-weight females matched on age and education to the AN and BN groups, and education only to the obese group, with no history of psychiatric or neurological illness, and no current clinical diagnosis. The group ranged in age from 18 to 40 years.

Both studies received ethical approval from the relevant hospital ethics committees, and all participants gave written informed consent.

## Measures

#### *Anthropometric measurements*

BMI (weight in kilograms/height in meters<sup>2</sup>) was measured to the nearest 0.1 Kg and 0.01 m.

#### *Decision making*

In a computerized version of the IGT (Bechara et al., 1994), participants are given a loan of \$2000 play money and are told to make choices that maximize their gains, with minimal instructions regarding the rules of the task. Over 100 trials, they choose between four decks of cards (A, B, C, D) some of which yield high immediate gain (\$100) but larger future losses (“disadvantageous decks”: A and B) and others that

yield lower immediate gain (\$50) but smaller future losses (“advantageous decks”: C and D). Every 10 cards from decks A and B gain \$1000, but deck A incurs five unpredictable punishments ranging from \$150 to \$350 bringing the total loss to \$1250, while deck B incurs one large punishment of \$1250. Every 10 cards from deck C or D amounts to a gain of \$500, but the punishments incurred are also smaller ranging from \$25 to \$75 in deck C with a total loss of \$250, and a single loss of \$250 in deck D. Decks A and B are disadvantageous because they cost the most in the long run, while decks C and D are advantageous because they result in an overall gain.

## Data Analysis

IGT performance between groups (comparison group, AN, BN, obese) was examined by a global outcome score (net score) and a net score for each consecutive block of 20 cards. Net scores are derived from the total number of cards chosen from the advantageous decks (C & D) minus the number of cards chosen from the disadvantageous decks (A & B). Lower scores represent poorer performance, and negative scores indicate a relative preference for choices from the disadvantageous decks. A total net score of <10 was established by Bechara, Dolan, Denburg, Hindes, Anderson, and Nathan (2001) as the threshold for impaired performance on the IGT, given the maximum net score achieved by VmPFC patients was <10. Analysis of IGT performance by blocks of 20 cards (blocks 1–5) provides an index of learning and strategy used by participants across the trials.

A mixed factorial analysis of covariance (ANCOVA) between the comparison, AN, BN, and obese groups compared differences on IGT global net scores, while a 4 (group)  $\times$  5 (block) repeated measures ANCOVA was carried out to examine group differences in block net scores. *Post hoc* analysis was conducted using the Sidak adjustment for multiple comparisons.

## RESULTS

### Sample Characteristics

The demographic profile of female participants is summarized in Table 1. There were significant differences between the groups in BMI ( $F(3,74) = 63.76; p < .001$ ), with the obese and BN groups significantly higher than the AN or comparison group, and in age ( $F(3,74) = 36.48; p < .001$ ) due to the older age of the obese group. Age was therefore entered as a covariate in all analyses. Ten obese participants (55%) had received bariatric surgery; as there were no significant differences between these participants and the non-bariatric surgery group on global net or block net scores, results are presented for the entire obese group.

### Global Net Scores

The mean net scores for each group are presented in Figure 1 (left panel). Controlling for age, there was a significant

difference on IGT global net score ( $F(3,73) = 11.06; p < .001$ ). *Post hoc* analysis revealed the comparison group was significantly different to the AN ( $p < .001$ ), BN ( $p < .001$ ), and obese groups ( $p = .004$ ); however, the three clinical groups were not significantly different from each other.

### Block Net Scores

The mean block net scores for each group are presented in Figure 1 (right panel). Controlling for age, there was a significant main effect for Group ( $F(3,73) = 12.06; p < .001$ ), but not for Block ( $F(3.49,254.97) = 0.26; p = .88$ ), and a significant Group  $\times$  Block interaction ( $F(10.48,254.97) = 3.23; p < .001$ ). *Post hoc* analysis revealed no significant group differences at blocks 1 and 2. However, the comparison group was significantly different to the AN, BN and obese groups at blocks 3 ( $p < .01$ ) and 4 ( $p < .01$ ), and from the AN ( $p < .05$ ) and obese ( $p < .05$ ) groups only at block 5. The three clinical groups were not significantly different from each other in any block.

### IGT Impairment

Sixty-one percent of AN, 77% of BN, and 72% of obese participants met the threshold (IGT<10) for impairment on the task (Bechara et al., 2001). By contrast, only 15% of the comparison group fell within this range.

## DISCUSSION

All three clinical groups were significantly impaired on the IGT relative to the comparison group. The majority of participants within the clinical groups reached the threshold for impairment on the IGT (although the clinical utility of this cutoff remains to be established). However, the clinical groups were not significantly different from each other, providing evidence of a possible shared inability to modulate reward and punishment in a long-term perspective among females with disordered eating. The IGT is a complex task that taps many aspects of decision making; several possible explanations for this shared decision making profile are discussed.

As the clinical groups did not perform within the negative range on the IGT (as in VmPFC patients), this indicates that they were somewhat sensitive to punishment; consequently their performance deficit may lie in faulty reward processing. The relatively flat learning curves of the three clinical groups (Figure 1) suggest that they could neither maximize an immediate reward (by choosing the disadvantageous decks) or a delayed reward (by choosing the advantageous decks). This may indicate learned helplessness (Hiroto & Seligman, 1975). An expectancy that the task is uncontrollable, with responding unrelated to reinforcement, may account for the flat performance of the clinical groups and may mimic their attempts to ameliorate pathological eating behaviors. For the obese group who showed a decline in performance over the last 20 trials (unlike the AN and BN

**Table 1.** Clinical Demographic Profile of Participants

	NC Mean (SD)	AN Mean (SD)	BN Mean (SD)	Obese Mean (SD)
Age	27.75 (6.99) <sup>a</sup>	29.09 (7.36) <sup>a</sup>	29.94 (6.41) <sup>a</sup>	52.11 (11.65) <sup>b</sup>
Years in education	12.15 (2.92)	12.96 (2.51)	11.65 (3.33)	14.00 (2.91)
BMI	21.55 (1.42) <sup>a</sup>	16.03 (2.04) <sup>b</sup>	31.87 (9.42) <sup>c</sup>	36.20 (5.04) <sup>c</sup>

BMI = Body Mass Index;

<sup>a,b,c</sup>Different superscript indicates significant differences between means.

groups who showed modest improvements from blocks 4 to 5), the task may mirror a likely long-standing pattern of failed weight loss attempts.


Unlike the comparison group who began to select more cards from the advantageous decks and less from the disadvantageous decks by the second block, the three clinical groups failed to learn across the task. As normal performance on the IGT appears to require *reversal learning*, in that the fixed card order induces an initial preference for the ultimately riskier decks (due to higher initial gains on these decks), this may provide an alternative explanation for the shared deficits in the female eating disordered groups. Imitating the choices on the IGT, individuals with pathological eating behavior have to learn to reverse behaviors that may once have been highly rewarding (e.g., dieting in AN, binging in BN, and overeating palatable foods in obesity) but ultimately bring higher costs in terms of deleterious physical and psychological outcomes. Therefore, reversal learning deficits present a viable explanation for the shared decision making profile and inability to learn in the three clinical groups on the IGT. In a shuffle variant of the IGT wherein the card order was changed, eliminating the need for reversal learning, Fellows and Farah (2005) found the performance of VmPFC patients improved to control levels. Future testing of the three clinical groups on this variant measure would inform this reversal learning hypothesis.

Impaired IGT performance among the three female pathological eating groups highlights the importance of adaptive decision making for successful self-regulation of eating be-

havior. However, following previous work with substance dependent individuals (Bechara & Damasio, 2002) use of both the original and variant IGT (in which the schedules of reward and punishment are reversed so that punishment is immediate and reward is delayed) in AN, BN, and obesity would allow differentiation of the nature of this deficit; specifically, whether it is attributable to hypersensitivity to reward, insensitivity to punishment, or a general “myopia” for future consequences (positive and negative) with behavior guided by immediate prospects.

### Limitations and Future Research

The obese group was significantly older than the AN, BN, and comparison group; however, this was controlled for statistically and research suggests that the capacity for adaptive self-regulation is a relatively stable disposition throughout the lifespan (Davis et al., 2010). Future research should take into account factors that were not controlled for here, such as illness duration, IQ, psychopathology, impulsivity, and medication use, which may impact upon IGT performance. Furthermore, the results are generalizable to females only and sample sizes were small, consequently replication with a larger group including males is desirable. Future research should also investigate whether interventions could improve IGT decision making capacity (in terms of the ability to modulate reward and punishment in a long-term perspective) and to assess subsequent impact on psychological and physical outcomes.



**Fig. 1.** (Left Panel) Mean net scores [(C&D – A&B)] for each group. (Right Panel) Mean net scores [(C&D – A&B)] for blocks 1 – 5 for each group. Positive scores reflect advantageous performance while negative scores reflect disadvantageous performance. Errors bars are standard errors.

## CONCLUSION

Females with AN, BN, and obesity were significantly impaired on the IGT with reference to a comparison group on both overall task performance and learning across the task. The three clinical groups did not differ significantly from each other, indicating that IGT performance may represent a common neuropsychological correlate of the patients' deficient eating behavior.

## ACKNOWLEDGMENTS

The information in this manuscript, and the manuscript itself, has never been published electronically or in print. This research was supported by the Irish Research Council for the Humanities and Social Sciences (IRCHSS).

## REFERENCES

- American Psychiatric Association. (2000). *Diagnostic and statistical manual of mental disorders: DSM-IV-TR* (4th ed.). Washington, DC: American Psychiatric Association.
- Bechara, A., & Damasio, H. (2002). Decision-making and addiction (I): Impaired activation of somatic states in substance dependent individuals when pondering decisions with negative future consequences. *Neuropsychologia*, *40*, 1675–1689.
- Bechara, A., Damasio, A.R., Damasio, H., & Anderson, S.W. (1994). Insensitivity to future consequences following damage to human prefrontal cortex. *Cognition*, *50*, 7–15.
- Bechara, A., Dolan, S., Denburg, N., Hindes, A., Anderson, S.W., & Nathan, P.E. (2001). Decision-making deficits, linked to a dysfunctional ventromedial prefrontal cortex, revealed in alcohol and stimulant abusers. *Neuropsychologia*, *39*, 376–389.
- Bechara, A., Tranel, D., Damasio, H., & Damasio, A.R. (1996). Failure to respond autonomically to anticipated future outcomes following damage to prefrontal cortex. *Cerebral Cortex*, *6*, 215–225.
- Boeka, A.G., & Lokkenz, K.L. (2006). The Iowa gambling task as a measure of decision making in women with bulimia nervosa. *Journal of the International Neuropsychological Society*, *12*, 741–745.
- Bosanac, P., Kurlender, S., Stojanovska, L., Hallam, K., Norman, T., McGrath, C., et al. (2007). Neuropsychological study of underweight and “weight-recovered” anorexia nervosa compared with bulimia nervosa and normal controls. *International Journal of Eating Disorders*, *40*, 613–621.
- Brand, M., Franke-Sievert, C., Jacoby, G.E., Markowitsch, H.J., & Tuschen-Caffier, B. (2007). Neuropsychological correlates of decision making in patients with bulimia nervosa. *Neuropsychology*, *21*, 742–750.
- Cavedini, P., Bassi, T., Ubbiali, A., Casolari, A., Giordani, S., Zorzi, C., et al. (2004). Neuropsychological investigation of decision-making in anorexia nervosa. *Psychiatry Research*, *127*, 259–266.
- Cavedini, P., Zorzi, C., Bassi, T., Gorini, A., Baraldi, C., Ubbiali, A., et al. (2006). Decision-making functioning as a predictor of treatment outcome in anorexia nervosa. *Psychiatry Research*, *145*, 179–187.
- Damasio, A.R. (1994). *Descartes error: Emotion, reason and the human brain*. London: Picador.
- Davis, C., Levitan, R., Muglia, P., Bewell, C., & Kennedy, J. (2004). Decision-making deficits and overeating: A risk model for obesity. *Obesity Research*, *12*, 929–935.
- Davis, C., Patte, K., Curtis, C., & Reid, C. (2010). Immediate pleasures and future consequences. A neuropsychological study of binge eating and obesity. *Appetite*, *54*, 208–213.
- Dunn, B.D., Dalgleish, T., & Lawrence, A.D. (2006). The somatic marker hypothesis: A critical evaluation. *Neuroscience & Biobehavioral Reviews*, *30*, 239–271.
- Fellows, L.K., & Farah, M.J. (2005). Different underlying impairments in decision-making following ventromedial and dorsolateral frontal lobe damage in humans. *Cerebral Cortex*, *15*, 58–63.
- Hiroto, D.S., & Seligman, M.E. (1975). Generality of learned helplessness in man. *Journal of Personality and Social Psychology*, *31*, 311–327.
- Liao, P.-C., Uher, R., Lawrence, N., Treasure, J., Schmidt, U., Campbell, I.C., et al. (2009). An examination of decision making in bulimia nervosa. *Journal of Clinical and Experimental Neuropsychology*, *31*, 455–461.
- Pignatti, R., Bertella, L., Albani, G., Mauro, A., Molinari, E., & Semenza, C. (2006). Decision-making in obesity: A study using the Gambling Task. *Eating and Weight Disorders*, *11*, 126–132.
- Tchanturia, K., Liao, P.-C., Uher, R., Lawrence, N., Treasure, J., & Campbell, I.C. (2007). An investigation of decision making in anorexia nervosa using the Iowa Gambling Task and skin conductance measurements. *Journal of the International Neuropsychological Society*, *13*, 635–641.