







FIRST ANALYSIS OF THE EFFECT OF THE JOVICENTRIC DECLINATION OF THE EARTH ON THE OBSERVATION OF JOVIAN DECAMETRIC RADIO EMISSIONS

H. R. P. Jácome^{1*} , M. S. Marques² , P. Zarka^{3,4} ,
E. Echer¹ , L. Lamy^{3,4,5} , and C. K. Louis⁶ 

*Corresponding author: hadassajacome@gmail.com

Citation:

Jácome et al., 2023, First analysis of the effect of the Jovicentric declination of the Earth on the observation of Jovian decametric radio emissions, in *Planetary, Solar and Heliospheric Radio Emissions IX*, edited by C. K. Louis, C. M. Jackman, G. Fischer, A. H. Sulaiman, P. Zucca, published by DIAS, TCD, pp. 241–249, doi: 10.25546/104049

Abstract

Through analysis of the extensive catalog of the Nançay Decameter Array, from 1978 to 2020, we demonstrate that the effect of the variation of the Jovicentric declination of the Earth on the visibility of Jovian decametric radio emissions by ground-based instruments is combined with the effects of variation of the Earth-Jupiter distance and of Jupiter's elongation. Therefore, these superimposed effects must be considered and removed for the study of the pure effect of the declination on the emissions' visibility.

¹ DIHPA, CGCE, Instituto Nacional de Pesquisas Espaciais, São José dos Campos, Brazil

² Departamento de Geofísica, Universidade Federal do Rio Grande do Norte, Natal, Brazil

³ LESIA, Observatoire de Paris, Université PSL, CNRS, Sorbonne Université, Université de Paris, Meudon, France

⁴ Observatoire de Radioastronomie de Nançay, Observatoire de Paris, Université PSL, CNRS, Univ. Orléans, Nançay, France

⁵ Aix Marseille Université, CNRS, CNES, LAM, Marseille, France

⁶ School of Cosmic Physics, DIAS Dunsink Observatory, Dublin Institute for Advanced Studies, Dublin 15, Ireland

1 Introduction

The Jovian decametric (DAM) radio emissions are the only known type of non-thermal planetary radiation that can be detected by ground-based radio instruments because of their high frequencies, up to 40 MHz, that overcome the cut-off frequency of the terrestrial ionosphere. For this reason, the Jovian DAM emissions were the first clue of the existence of a magnetic field and magnetosphere at Jupiter, and have been continuously observed by ground-based instruments for decades. One factor that is long known to affect the visibility of the Jovian DAM emissions by ground-based instruments, besides the cut-off frequency of the terrestrial ionosphere limiting the minimum observed frequency, is the variation of the Jovicentric declination of the Earth (D_E), i.e., the sub-Earth Jovicentric latitude. It has been observed that, as D_E varies, the distribution of each component of the Jovian DAM emissions as a function of the longitude and of Io orbital phase changes, for instance, in amplitude and in width. This could result from the position of the radiation beaming cones, (cyclotron maser instability cones Wu & Lee, 1979) relative to the observer that becomes more or less visible as the declination changes (Carr et al., 1970; Boudjada & Leblanc, 1992; Leblanc et al., 1993; Garcia, 1996; Imai et al., 2011, and references therein). Advances in the study of Jovian DAM radiation generation and in simulations of their visibility have improved the understanding of the observation constraints and, consequently, of the geometry of the radiation beaming cones, which are most probably oblate cones centered in Jovian magnetic field lines (Galopeau & Boudjada, 2016; Louis et al., 2017). In this context, the comprehension of the declination effect on the Jovian DAM emission visibility could contribute to validate and possibly improve the beaming cone morphology.

However, although some aspects of the effect of the variation of D_E have been observed and studied, a real and clear relation between it and the visibility of Jovian DAM emissions is still an open question that might be answered through a long-term, multidecadal study of the emissions variability with D_E . The radio observation of Jupiter by the Nançay Decameter Array (NDA) since 1978 provides an extensive database of Jovian DAM emissions (Lamy et al., 2017), which in turn allows in-depth studies of the Jovian DAM components (Zarka et al., 2017; Marques et al., 2017; Zarka et al., 2018; Jácome et al., 2022) and enables the study of their long-term variability.

A surprising behavior of the daily observations of Jupiter by the NDA and of the detected Jovian DAM emissions is that the majority of them are distributed in clusters around specific values of D_E , such as -3° , -1.5° , 0° , 2° and 3.5° , with just a relatively few cases occurring between these clusters, as shown in the histogram of Figure 1a. This type of distribution is even clearer for the emissions, which seems to indicate that the visibility of the emissions is favoured when the declination is around those values. However, more intriguing than that is the conservation of the clustered distribution when analyzing the ratio of the number of emissions to the number of observations for each bin of declination, showed in the panel *b*. It suggests that the occurrence of observations of Jupiter with no emission detection is less frequent around the declination values of -3° , -1.5° , 0° , 2° and 3.5° . The same inference may be deduced from the distribution of the emissions' occurrence probability, shown in Figure 1d. The probability was calculated from the ratio of the sum of the duration of the emissions to the sum of the duration of the observations

for each 0.25° of declination. The distribution of the total duration of the emissions and of the observations found in each 0.25° bin is shown in panel *c*.

If one considers the variation of the declination in time, shown in Figure 2, it can be noted that the Earth spends more time in D_E around -3° , -1.5° , 0° , 2° and 3.5° , which could explain the higher amount of observations and detected emissions around these values. But what about the detection probability? We would expect it to present an approximately flat distribution in declination if detection depended only on the amount of time that the Earth spends around declination values more favourable to Jovian emissions' observation. As we still observe a modulation vs. D_E of the occurrence probability, this means that other factors must also affect the detection of Jovian DAM emissions.




Figure 1: (a) Number of observations (black) and detected emissions (red) found in intervals of 0.25° of the jovicentric declination of the Earth. (b) Ratio of the number of emissions to the number of observations in each bin of the histogram in panel a. (c) The sum of duration, in hours, of the observations (black) and of the emissions (red) found in each 0.25° bin. (d) Emissions' occurrence probability (%), calculated from the ratio of the sum of duration of the emissions to the sum of duration of the observations for each interval of declination.




Figure 2: Daily variation of the Jovicentric declination of the Earth, from Jan. 3th, 1978 to Dec. 31st, 2020.

In this work, we identify two other factors of the relative motion between the Earth and Jupiter that also affect the visibility of the Jovian DAM emissions and are combined

with the D_E effect: the variation of the distance between the Earth and Jupiter and the variation of Jupiter's elongation, both sketched in Figure 3.

2 Method

We use the extensive digital catalog of the NDA, which comprises all the observations of Jupiter by the array from January 3, 1978 to December 31, 2020, with all the detected emissions in this period (Lamy et al., 2017; Marques et al., 2017). We have plotted the distributions of all the observations of Jupiter by the NDA, all the Jovian DAM emissions catalogued and the occurrence probability of the emissions as a function of D_E versus the Earth-Jupiter distance (Figure 4) and of D_E versus Jupiter's elongation (Figure 5).

The Earth-Jupiter distance, which varies from ~ 4.0 AU to 6.5 AU, can affect the intensity of the detected emissions, with the less intense ones being detected only at shortest distances. Jupiter's elongation affects the minimum frequency of the detected emissions, with emissions with lower frequency (e.g., freq. < 25 MHz) being detected only when the planets are close to opposition (i.e., when Jupiter is observed in the Earth's night-side sky), due to the increased radio interference in the day side that limit the visibility of low frequencies.

Figure 3 shows a sketch of the Earth and Jupiter and the distance (R) and Jupiter's elongation (θ). R is the distance between the planets at the meridian transit of Jupiter for each observation, given in the NDA catalog. θ is the angle between the Sun and Jupiter, observed from the Earth, with $\theta \rightarrow 0^\circ$ indicating that the planets are close to conjunction; and $\theta \rightarrow \pm 180^\circ$, that the planets are close to opposition. Jupiter's elongation was collected for each day from January 1st, 1978 to December 31, 2020, from NASA's Horizons System¹.




Figure 3: Sketch (not to scale) of the Earth (in blue) and Jupiter (in orange) on their orbits, with the distance (R) and Jupiter's elongation (θ). When $\theta = 0^\circ$, the planets are in conjunction, and when $\theta = \pm 180^\circ$, they are in opposition.

3 Results

Figure 4 shows distributions of all the observations of Jupiter by the NDA, from 1978 to 2020 (panel *a*), of all the detected Jovian DAM emissions (panel *b*) and of the occurrence

¹<https://ssd.jpl.nasa.gov/horizons/app.html>

probability of those emissions (panel *c*) as a function of D_E and of the Earth-Jupiter distance. It is observed that, for distances shorter than 5.5 AU, the observations occur around D_E values of -3° , -1.5° , 0° , 2° and 3.5° , which explains why the majority of the observations accumulate around those values. However, for longer distances, observations occur over the entire D_E range. We also note that the observations are quite homogeneously distributed in distance, with only a smooth increase in number (panel *a*) around the shortest (~ 4.25 AU) and the longest (~ 6.25 AU) distances.




Figure 4: Distributions of all the observations of Jupiter (*a*) by the Nançay Decameter Array, from 1978 to 2020; of all the detected Jovian DAM emissions (*b*); and of the emissions' occurrence probability (*c*), as a function of the jovicentric declination of the Earth and of the Earth-Jupiter distance.

The distribution of the emissions (panel *b* of Figure 4), on the other hand, shows that although emissions are detected over the entire distance range, they accumulate at the shortest distances, which indicates that the emissions detection is favoured by the shortest distance between the Earth and Jupiter. As a consequence, the emissions' occurrence

probability (panel *c*) is higher at the shortest distances, as observed in Figure 1.

In summary, although Jupiter is observed by the NDA over the entire Earth-Jupiter distance range, most of the emissions are detected when this distance is shortest, which always coincides with D_E values around -3° , -1.5° , 0° , 2° and 3.5° .

Figure 5 shows distributions of all the observations of Jupiter by the NDA, from 1978 to 2020 (panel *a*), of all the detected Jovian DAM emissions (panel *b*) and of the occurrence probability of those emissions (panel *c*) as a function of D_E and of Jupiter's elongation. In panel *a*, we see that the observations are homogeneously distributed over the entire range of the elongation angle. The emissions visibility (panel *b*), however, is favoured when the planets are close to opposition ($\theta \rightarrow \pm 180^\circ$). Otherwise, only a few emissions are detected, most probably the ones with the highest frequencies. As the planets opposition coincides with D_E values around -3° , -1.5° , 0° , 2° and 3.5° , this explains why the emissions and their higher occurrence probability accumulate around those values.

In summary, although Jupiter is observed by the NDA over the entire range of the planet's elongation, most of the emissions are detected when it is around opposition with the Earth, which always coincides with D_E values around -3° , -1.5° , 0° , 2° and 3.5° .

4 Conclusions and perspectives

We have presented an initial analysis of the effect of the variation of Jovicentric declination of the Earth on the detection of Jovian DAM emissions by ground-based instruments such as the Nançay Decameter Array. We have demonstrated that the detection of emissions is favoured when the planets are close to opposition ($\theta \rightarrow \pm 180^\circ$) and the distance between the planets is smallest (less than 5.5 AU), which both coincide with D_E around -3° , -1.5° , 0° , 2° and 3.5° . Therefore, the D_E effect is actually combined with the effect of the variation of the distance and of Jupiter's elongation. Our first conclusion is that the results of all past studies of the D_E effect (e.g., Barrow, 1981) are unreliable because the effects of the Earth-Jupiter distance and of Jupiter's elongation associated with it were not identified and thus not corrected. In order to study and understand the real declination effect, those other superimposed effects must be removed by adequate data selections.

For the next step, we intend to remove the effect of the distance and of Jupiter's elongation by selecting emissions whose observation is not limited by distance or radio interference, i.e., emissions that can be detected over the entire ranges of Earth-Jupiter distance (~ 4 – 6.5 AU) and of Jupiter's elongation (from opposition to conjunction). Then, we will study the pure D_E effect on the Jovian DAM emissions visibility.

The results will be compared to simulations of the emissions' dynamic spectra, with EXPRES (Louis et al., 2019), in order to interpret them.




Figure 5: Distributions of all the observations of Jupiter (a) by the Nançay Decameter Array, from 1978 to 2020; of all the detected Jovian DAM emissions (b); and of the emissions' occurrence probability (c), as a function of the jovicentric declination of the Earth and of Jupiter's elongation.

Acknowledgements

This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) - Finance Code 001. E. Echer would like to thank the Brazilian agencies CNPq (PQ-301883/2019-0) and FAPESP (2018/21657-1) for the research grants. C. K. Louis' work at DIAS is supported by the Science Foundation Ireland Grant 18/FRL/6199. The authors thank the Brazilian Ministry of Science, Technology and Innovation and the Brazilian Space Agency. The French co-authors acknowledge support from CNES and CNRS/INSU programs of planetology (PNP) and heliophysics (PNST).

References

- Barrow C. H., 1981, Latitudinal beaming and local time effects in the decametre-wave radiation from Jupiter observed at the Earth and from Voyager, *Astronomy and Astrophysics*, 101, 142
- Boudjada M. Y., Leblanc Y., 1992, The variability of Jovian decametric radiation from 1978 to 1988, *Advances in Space Research*, 12, 95
- Carr T. D., Smith A. G., Donovan F. F., Register H. I., 1970, The twelve-year periodicities of the decametric radiation of Jupiter, *Radio Science*, 5, 495
- Galopeau P. H. M., Boudjada M. Y., 2016, An oblate beaming cone for Io-controlled Jovian decameter emission, *Journal of Geophysical Research (Space Physics)*, 121, 3120
- Garcia L. N., 1996, Long-term periodicities in the Jovian decametric emission, *PhD thesis, University of Florida*
- Imai K., Garcia L., Reyes F., Imai M., Thieman J. R., 2011, A model of Jupiter's decametric radio emissions as a searchlight beam, *Planetary, Solar and Heliospheric Radio Emissions (PRE VII)*, pp 179–186
- Jácome H. R. P., Marques M. S., Zarka P., Echer E., Lamy L., Louis C. K., 2022, Search for Jovian decametric emission induced by Europa on the extensive Nançay Decameter Array catalog, *Astronomy & Astrophysics*, 665, A67
- Lamy L., Zarka P., Cecconi B., Klein L., Masson S., Denis L., Coffre A., Viou C., 2017, 1977-2017: 40 years of decametric observations of Jupiter and the Sun with the Nançay Decameter Array, in *Planetary Radio Emissions VIII*, eds Fischer, G. and Mann, G. and Panchenko, M. and Zarka, P., pp 455–466, doi:10.1553/PRE8s455
- Leblanc Y., Gerbault A., Denis L., Lecacheux A., 1993, A catalogue of Jovian decametric radio observations from January 1988 to December 1990, *Astronomy and Astrophysics Supplement Series*, 98, 529
- Louis C. K., et al., 2017, Io-Jupiter decametric arcs observed by Juno/Waves compared to ExPRES simulations, *Geophysical Research Letters*, 44, 9225
- Louis C. K., Hess S. L. G., Cecconi B., Zarka P., Lamy L., Aicardi S., Loh A., 2019, ExPRES: an Exoplanetary and Planetary Radio Emissions Simulator, *Astronomy & Astrophysics*, 627, A30
- Marques M. S., Zarka P., Echer E., Ryabov V. B., Alves M. V., Denis L., Coffre A., 2017, Statistical analysis of 26 yr of observations of decametric radio emissions from Jupiter, *Astronomy & Astrophysics*, 604, A17
- Wu C., Lee L., 1979, A theory of the terrestrial kilometric radiation, *The Astrophysical Journal*, 230, 621

Zarka P., Marques M. S., Louis C., Ryabov V. B., Lamy L., Echer E., Cecconi B., 2017, Radio emission from satellite-Jupiter interactions (especially Ganymede), *in Planetary Radio Emissions VIII*, eds Fischer, G. and Mann, G. and Panchenko, M. and Zarka, P., pp 45–58, doi:10.1553/PRE8s45

Zarka P., Marques M. S., Louis C., Ryabov V. B., Lamy L., Echer E., Cecconi B., 2018, Jupiter radio emission induced by Ganymede and consequences for the radio detection of exoplanets, *Astronomy & Astrophysics*, 618, A84