

National Parks and Wildlife Service

Conservation Objectives Series

Hugginstown Fen SAC 000404



An Roinn Cultúir,
Oidhreachta agus Gaeltachta
Department of Culture,
Heritage and the Gaeltacht

**National Parks and Wildlife Service,
Department of Culture, Heritage and the Gaeltacht,
90 King Street North, Dublin 7, D07 N7CV, Ireland.**

**Web: www.npws.ie
E-mail: nature.conservation@chg.gov.ie**

Citation:

**NPWS (2019) Conservation Objectives: Hugginstown Fen SAC 000404. Version 1.
National Parks and Wildlife Service, Department of Culture, Heritage and the
Gaeltacht.**

**Series Editor: Rebecca Jeffrey
ISSN 2009-4086**

Introduction

The overall aim of the Habitats Directive is to maintain or restore the favourable conservation status of habitats and species of community interest. These habitats and species are listed in the Habitats and Birds Directives and Special Areas of Conservation and Special Protection Areas are designated to afford protection to the most vulnerable of them. These two designations are collectively known as the Natura 2000 network.

European and national legislation places a collective obligation on Ireland and its citizens to maintain habitats and species in the Natura 2000 network at favourable conservation condition. The Government and its agencies are responsible for the implementation and enforcement of regulations that will ensure the ecological integrity of these sites.

A site-specific conservation objective aims to define favourable conservation condition for a particular habitat or species at that site.

The maintenance of habitats and species within Natura 2000 sites at favourable conservation condition will contribute to the overall maintenance of favourable conservation status of those habitats and species at a national level.

Favourable conservation status of a habitat is achieved when:

- its natural range, and area it covers within that range, are stable or increasing, and
- the specific structure and functions which are necessary for its long-term maintenance exist and are likely to continue to exist for the foreseeable future, and
- the conservation status of its typical species is favourable.

The favourable conservation status of a species is achieved when:

- population dynamics data on the species concerned indicate that it is maintaining itself on a long-term basis as a viable component of its natural habitats, and
- the natural range of the species is neither being reduced nor is likely to be reduced for the foreseeable future, and
- there is, and will probably continue to be, a sufficiently large habitat to maintain its populations on a long-term basis.

Notes/Guidelines:

1. The targets given in these conservation objectives are based on best available information at the time of writing. As more information becomes available, targets for attributes may change. These will be updated periodically, as necessary.
2. An appropriate assessment based on these conservation objectives will remain valid even if the targets are subsequently updated, providing they were the most recent objectives available when the assessment was carried out. It is essential that the date and version are included when objectives are cited.
3. Assessments cannot consider an attribute in isolation from the others listed for that habitat or species, or for other habitats and species listed for that site. A plan or project with an apparently small impact on one attribute may have a significant impact on another.
4. Please note that the maps included in this document do not necessarily show the entire extent of the habitats and species for which the site is listed. This should be borne in mind when appropriate assessments are being carried out.
5. When using these objectives, it is essential that the relevant backing/supporting documents are consulted, particularly where instructed in the targets or notes for a particular attribute.

Qualifying Interests

** indicates a priority habitat under the Habitats Directive*

000404 Hugginstown Fen SAC

7230 Alkaline fens

Supporting documents, relevant reports & publications

Supporting documents, NPWS reports and publications are available for download from: www.npws.ie/Publications

NPWS Documents

Year :	1972
Title :	A Preliminary Report on Areas of Scientific Interest in County Kilkenny
Author :	Young, R.
Series :	Unpublished Report
Year :	2009
Title :	Ireland Red List No. 2: Non-marine molluscs
Author :	Byrne, A.; Moorkens, E.A.; Anderson, R.; Killeen, I.J.; Regan, E.C.
Series :	Ireland Red List series, NPWS
Year :	2010
Title :	Ireland Red List No. 4: Butterflies
Author :	Regan, E.C.; Nelson, B.; Aldwell, B.; Bertrand, C.; Bond, K.; Harding, J.; Nash, D.; Nixon, D.; Wilson, C.J.
Series :	Ireland Red List series, NPWS
Year :	2011
Title :	Ireland Red List No. 6: Damselflies and Dragonflies (Odonata)
Author :	Nelson, B.; Ronayne, C.; Thompson, R.
Series :	Ireland Red Lists series, NPWS
Year :	2012
Title :	Ireland Red List No. 8: Bryophytes
Author :	Lockhart, N.; Hodgetts, N.; Holyoak, D.
Series :	Ireland Red List series, NPWS
Year :	2013
Title :	The status of EU protected habitats and species in Ireland. Volume 2. Habitats assessments
Author :	NPWS
Series :	Conservation assessments
Year :	2014
Title :	Guidelines for a national survey and conservation assessment of upland vegetation and habitats in Ireland, Version 2.0
Author :	Perrin, P.M.; Barron, S.J.; Roche, J.R.; O'Hanrahan, B.
Series :	Irish Wildlife Manuals, No. 79
Year :	2016
Title :	Ireland Red List No. 10: Vascular Plants
Author :	Wyse Jackson, M.; FitzPatrick, Ú.; Cole, E.; Jebb, M.; McFerran, D.; Sheehy Skeffington, M.; Wright, M.
Series :	Ireland Red List Series, NPWS
Year :	2018
Title :	Backing document – National Conservation Status Assessments (NCAs) for three fen habitat types: 7140 – Transition mires and quaking bogs, 7210 – Calcareous fens with <i>Cladium mariscus</i> and species of <i>Caricion davallianae</i> , 7230 – Alkaline fens
Author :	Long, M.P.; Crowe, O.; Kimberley, S.; Denyer, J.
Series :	Unpublished report to NPWS
Year :	in prep.
Title :	The Status of EU Protected Habitats and Species in Ireland (2013-2018). Habitat Assessments
Author :	NPWS
Series :	Conservation assessments

Other References

- Year :** 2004
Title : Common Standards Monitoring guidance for lowland wetland habitats
Author : JNCC
Series : Joint Nature Conservation Committee, Peterborough
-
- Year :** 2011
Title : Review and revision of empirical critical loads and dose-response relationships. Proceedings of an expert workshop, Noordwijkerhout, 23-25 June 2010
Author : Bobbink, R.; Hettelingh, J.P.
Series : RIVM report 680359002, Coordination Centre for Effects, National Institute for Public Health and the Environment (RIVM)
-
- Year :** 2018
Title : Irish Vegetation Classification: Technical Progress Report No. 4
Author : Perrin, P.
Series : Report submitted to National Biodiversity Data Centre
-

Conservation Objectives for : Hugginstown Fen SAC [000404]


7230 Alkaline fens

To maintain the favourable conservation condition of Alkaline fens in Hugginstown Fen SAC, which is defined by the following list of attributes and targets:

Attribute	Measure	Target	Notes
Habitat area	Hectares	Area stable or increasing, subject to natural processes	Alkaline fen has not been mapped in detail for Hugginstown Fen SAC and thus the total current area of the qualifying habitat in the SAC is unknown. According to Young (1972), the SAC contains one of the most important examples of the habitat in the region. The habitat is partly developed on mats of floating vegetation and occurs in association with common reed (<i>Phragmites australis</i>) swamp, which dominates the northern third of the SAC, bulrush (<i>Typha latifolia</i>) beds, small areas of transition mire, willow (<i>Salix</i> spp.) scrub and wet grassland (NPWS internal files)
Habitat distribution	Occurrence	No decline, subject to natural processes	See the notes for Habitat area above
Ecosystem function: soil nutrients	Soil pH and appropriate nutrient levels at a representative number of monitoring stops	Maintain soil pH and nutrient status within natural ranges	Relevant nutrients and their natural ranges are yet to be defined. However, nitrogen deposition is noted as being relevant to this habitat in NPWS (2013). See also Bobbink and Hettelingh (2011)
Ecosystem function: peat formation	Percentage cover of peat-forming vegetation and water table levels	Maintain active peat formation, where appropriate	In order for peat to form, water levels need to be slightly below or above the soil surface for c.90% of the time
Ecosystem function: hydrology - groundwater levels	Water levels (centimetres); duration of levels; hydraulic gradients	Maintain, or where necessary restore, appropriate natural hydrological regimes necessary to support the natural structure and functioning of the habitat	Fen habitats require high groundwater levels (i.e. water levels at or above the ground surface) for a large proportion of the calendar year (i.e. duration of mean groundwater level). Fen groundwater levels are controlled by regional groundwater levels in the contributing catchment area (which sustain the hydraulic gradients of the fen groundwater table). Regional abstraction of groundwater may affect fen groundwater levels. In this SAC, the swamp and fen habitats that dominate the SAC occupy a narrow low-lying basin on limestone glacial till overlying and surrounded by Old Red Sandstone. There are two catchments in the SAC, the Derrylackey River catchment which covers most of the fen and the Little Arrigle River catchment, which drains into the River Nore, in the very north of the fen. The fen is fed with iron-rich springs, which discharge along the eastern margin of the fen, and may also receive water from surrounding land (NPWS internal files)
Ecosystem function: hydrology - surface water flow	Drain density and form	Maintain, or where necessary restore, as close as possible to natural or semi-natural, drainage conditions	Drainage, either within or surrounding the fen habitat, can result in the drawdown of the alkaline fen groundwater table. The depth, geometry and density of drainage (hydromorphology) will indicate the scale and impact on fen hydrology. Drainage can result in loss of characteristic species and transition to drier habitats. A drain in the southern part of the SAC has impacted on the surface water and groundwater recharge in the southern fen, this drains the fen to the south into the Derrylackey River (NPWS internal files)
Ecosystem function: water quality	Water chemistry measures	Maintain appropriate water quality, particularly pH and nutrient levels, to support the natural structure and functioning of the habitat	Fens receive natural levels of nutrients (e.g. iron, magnesium and calcium) from water sources. However, they are generally poor in nitrogen and phosphorus, with the latter tending to be the limiting nutrient under natural conditions. Water supply should be also relatively calcium-rich. In this SAC, the chemistry of the groundwater indicates that it is alkaline and of calcium/magnesium bicarbonate water type. It has moderate mineral and nutrient content (NPWS internal files)

Community diversity	Abundance of variety of vegetation communities	Maintain variety of vegetation communities, subject to natural processes	The entire diversity of alkaline fen vegetation communities present in the SAC is currently unknown. Information on vegetation communities associated with this habitat in the uplands is presented in Perrin et al. (2014). See also the Irish Vegetation Classification (Perrin, 2018; www.biodiversityireland.ie/projects/national-vegetation-database/irish-vegetation-classification)
Vegetation composition: brown mosses	Percentage cover at a representative number of 2m x 2m monitoring stops	Maintain adequate cover of typical brown moss species	For lists of typical brown moss species, including high quality indicator species, see the 2013-2018 Article 17 conservation status assessment for alkaline fens (NPWS, in prep.) and the fen habitats supporting document (Long et al., 2018). In this SAC, the surface of the fen is covered in a carpet of mosses, including <i>Calliergonella cuspidata</i> (NPWS internal files)
Vegetation composition: typical vascular plants	Percentage cover at a representative number of 2m x 2m monitoring stops	Maintain adequate cover of typical vascular plant species	For lists of typical vascular plant species, including high quality indicator species, see the Article 17 conservation status assessment for alkaline fens (NPWS, in prep.) and the fen habitats supporting document (Long et al., 2018). See also Perrin et al. (2014) and JNCC (2004). In this SAC, the habitat supports a range of sedges including the typical species long-stalked yellow-sedge (<i>Carex lepidocarpa</i>), common sedge (<i>C. nigra</i>), carnation sedge (<i>C. panicea</i>), bottle sedge (<i>C. rostrata</i>) and black bog-rush (<i>Schoenus nigricans</i>), as well as other typical species including jointed rush (<i>Juncus articulatus</i>), water mint (<i>Mentha aquatica</i>), marsh pennywort (<i>Hydrocotyle vulgaris</i>), common marsh-bedstraw (<i>Galium palustre</i>), lesser spearwort (<i>Ranunculus flammula</i>) and the high quality indicators marsh helleborine (<i>Epipactis palustris</i>) and other orchids (<i>Dactylorhiza</i> spp.) (NPWS internal files)
Vegetation composition: native negative indicator species	Percentage cover at a representative number of 2m x 2m monitoring stops	Cover of native negative indicator species at insignificant levels	Negative indicators include species not characteristic of the habitat and species indicative of undesirable activities such as overgrazing, undergrazing, nutrient enrichment, agricultural improvement or impacts on hydrology. Native negative indicators may include graminoids such as reed canary-grass (<i>Phalaris arundinacea</i>) and reed sweet-grass (<i>Glyceria maxima</i>), tall herbs such as great willowherb (<i>Epilobium hirsutum</i>), bracken (<i>Pteridium aquilinum</i>), bramble (<i>Rubus fruticosus</i>) and common nettle (<i>Urtica dioica</i>), and bryophytes such as <i>Brachythecium rutabulum</i> and <i>Kindbergia praelonga</i>
Vegetation composition: non-native species	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of non-native species less than 1%	Attribute and target based on Perrin et al. (2014). Non-native species can be invasive and have deleterious effects on native vegetation. A low target is set as non-native species can spread rapidly and are most easily dealt with when still at lower abundances
Vegetation composition: native trees and shrubs	Percentage cover in local vicinity of a representative number of monitoring stops	Cover of scattered native trees and shrubs less than 10%	Attribute and target based on Perrin et al. (2014). Scrub and trees will tend to invade if fen conditions become drier
Vegetation composition: soft rush and common reed cover	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of soft rush (<i>Juncus effusus</i>) and common reed (<i>Phragmites australis</i>) less than 10%	Attribute and target based on Perrin et al. (2014)
Vegetation structure: litter	Percentage cover in local vicinity of a representative number of monitoring stops	Total cover of litter not more than 25%	Attribute and target based on JNCC (2004). More than 25% litter cover may indicate insufficient removal of biomass by grazing and/or undesirable water table levels

Physical structure: disturbed bare ground	Percentage cover at, and in local vicinity of, a representative number of 2m x 2m monitoring stops	Cover of disturbed bare ground not more than 10%	Attribute and target based on Perrin et al. (2014). While grazing may be appropriate in this habitat, excessive areas of disturbed bare ground may develop due to unsuitable grazing regimes. Disturbance can include hoof marks, wallows, human footprints, vehicle and machinery tracks. Excessive disturbance can result in loss of characteristic species and presage erosion for peatlands
Physical structure: tufa formations	Percentage cover in local vicinity of a representative number of monitoring stops	Disturbed proportion of vegetation cover where tufa is present is less than 1%	Attribute and target based on Perrin et al. (2014)
Indicators of local distinctiveness	Occurrence and population size	No decline in distribution or population sizes of rare, threatened or scarce species associated with the habitat; maintain features of local distinctiveness, subject to natural processes	This includes species on the Flora (Protection) Order, 2015 and/or Red Lists (Byrne et al., 2009; Regan et al., 2010; Lockhart et al., 2012; Wyse Jackson et al., 2016, etc.). The fen in the SAC is fed by a number of calcareous and iron-rich springs. The Near Threatened tubular water-dropwort (<i>Oenanthe fistulosa</i>) (Wyse Jackson et al., 2016) has been recorded in the habitat in the SAC (NPWS internal files). Some rare insects have been recorded in the SAC, notably the Near Threatened damselfly, robust spreadwing (<i>Lestes dryas</i>) (Nelson et al., 2011) and the hoverfly <i>Parhelophilus consimilis</i> (NPWS internal files)



Legend
 Hugginstown Fen SAC 000404

An Roinn Cultúir,
 Oidhreacht agus Gaeltachta
 Department of Culture,
 Heritage and the Gaeltacht

**MAP 1:
 HUGGINSTOWN FEN SAC
 CONSERVATION OBJECTIVES
 SAC DESIGNATION**
 Map to be read in conjunction with the NPWS Conservation Objectives Document.

**SITE CODE:
 SAC 000404; version 3.01. CO. KILKENNY**

0 180 360 540 720 Meters

The mapped boundaries are of an indicative and general nature only. Boundaries of designated areas are subject to revision.
 Ordnance Survey of Ireland Licence No EN 0059216. © Ordnance Survey of Ireland Government of Ireland

Níl sna teorainneacha ar na léarscáileanna ach nod garshuíomhach ginearálta. Féadfar athbhreithnithe a déanamh ar theorainneacha na gceantar comharthaite. Suirbhéarachta Ordonáis na hÉireann Ceadúnas Uimh EN 0059216. © Suirbhéarachta Ordonáis na hÉireann Rialtas na hÉireann

N
Map Version 1
Date: Jan 2019