

The Syncope-Falls Index (SYFI): A tool for predicting risk of syncope and complex falls in the older adult based on cumulative health deficits

Authors:

Neasa Fitzpatrick^{1, 2}, Roman Romero-Ortuno^{1,2,3}

Affiliated Institutions

1. MedEL Directorate, St. James's Hospital, James St, Dublin 8, Ireland
2. Discipline of Medical Gerontology, School of Medicine, Trinity College Dublin, Dublin 2, Ireland
3. Global Brain Health Institute, Trinity College Dublin, Ireland

Corresponding Author

Neasa Fitzpatrick, fitzpane@tcd.ie

Abstract

Background

Syncope is aetiologically diverse and associated with adverse outcomes; in older people, there is clinical overlap with complex falls presentations (i.e. recurrent, unexplained, and/or injurious).

Aim:

To formulate an index to predict future risk of syncope and falls in the Irish Longitudinal Study on Ageing (TILDA).

Design/Methods:

Using the frailty index methodology, we selected, from TILDA Wave 1 (2010), 40 deficits that might increase risk of syncope and falls. This syncope-falls index (SYFI) was applied to TILDA Wave 1 participants aged 65 and over, who were divided into three risk groups (low, intermediate, high) based on SYFI tertiles. Multivariate logistic regression models were used to investigate, controlling for age and sex, how SYFI groups predicted incident syncope, complex falls and simple falls occurring up to Wave 4 of the study (2016).

Results:

At Wave 1, there were 3499 participants (mean age 73, 53% women). By Wave 4, of the remaining 2907 participants, 185 (6.4%) had reported new syncope, 1077 (37.0%) complex falls, and 218 (7.5%) simple falls. The risk of both syncope and complex falls increased along the SYFI groups (high risk group: OR 1.88 [1.26 – 2.80], $P = 0.002$ for syncope; 2.22 [1.82 – 2.72], $P < 0.001$ for complex falls). No significant relationship was identified between SYFI and simple falls.

Conclusion:

The 6-year incidences of falls and syncope were high in this cohort. SYFI could help identify older adults at risk of syncope and complex falls, and thus facilitate early referral to specialist clinics to improve outcomes.

Keywords: syncope; falls; longitudinal; frailty index; older people.

Introduction

Syncope refers to a transient loss of consciousness due to transient global cerebral hypoperfusion characterised by rapid onset, short duration, and spontaneous complete recovery.(1) In younger cohorts, reflex vasovagal syncope is the most common form and does not reflect underlying pathology across physiological systems. However, in older adults the etiology of syncope can be more complex (2), with multiple causes frequently identified for syncope in one individual(3, 4). In older people, the proportions of cardiac and orthostatic syncope are higher, and these types have a multiplicity of associated risk factors including pathologies across many different organ systems (1). Multi-cause syncope is associated with poorer outcomes, with 77% vs. 89% survival at 4 years in those with two or more causes identified for their syncope.(4) Additionally, syncope and falls have been linked to geriatric syndromes such as cognitive impairment, polypharmacy and frailty.(3, 5, 6) Given that frailty is defined as dysregulation in multiple physiological systems with increased risk of decompensation in the face of stressors (7), the associations between frailty and syncope, and frailty and falls, are biologically plausible in older adults.

In older people, there is a recognised clinical overlap between syncope and complex falls presentations (i.e. recurrent, unexplained and/or injurious).(3) The possible syncopal origin of complex falls in older people remains underappreciated in non-specialist settings. Alongside increased mortality, syncope and complex falls in the older adult can lead to premature disability, as well as reductions in confidence, functionality and quality of life, and increased risk of institutionalisation.(8-10) Early detection and specialist intervention in those at risk of syncope and complex falls has been shown to effectively reduce adverse outcomes.(11, 12) Hence, it is important to bring this to the attention of non-specialist clinicians by offering simple tools that may help identify such patients.

1
2
3 In 2001, Mitnitski et al. formulated a Frailty Index (FI) methodology by which the cumulative
4 burden of health deficits acts as a surrogate measure of frailty.(13) A FI can be constructed
5 on any suitable health database provided that some requirements are met.(14) Most FIs
6 have concentrated on the prediction of mortality (15) but it is possible to consider more
7 specific health outcomes when selecting deficits for an FI.(16) Building on the clinical utility
8 of FIs in predicting outcomes in the older population, we formulated an FI designed
9 specifically to predict risk of syncope and falls and assess its predictive value on a
10 population-based cohort.
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Methods

Sample

The study utilised four waves of data from The Irish Longitudinal Study on Ageing (TILDA).(17-20) Authors accessed TILDA data via the Irish Social Science Data Archive (ISSDA) – www.ucd.ie/issda from July to December 2020. Full details of the recruitment and data collection process have previously been published.(21, 22)

The first wave of data (Wave 1) was collected between October 2009 and February 2011 and included participants aged 50 or more, and some younger spouses of the latter. Subsequent data was collected approximately 2-yearly and Wave 4 took place between January to December 2016. There was approximately 6-10% reduction in sample size between waves due to attrition, with death of participants between waves contributing another 2-3% reduction. For the purpose of our analyses, we included only those who were aged 65 or older at time of first interview.

Measures

Outcomes

Syncope – participants were asked in each wave if they had a faint or blackout in the past 12 months (yes/no). A case of syncope was defined as having answered yes to this question at least once at Wave 2, 3 or 4.

Falls – participants were asked in each wave if they had experienced a fall in the past 12 months (yes/no). A case of fall was defined as having answered yes to this question at least once at Wave 2, 3 or 4. Participants were asked the following questions to further characterise their falls:

1. How many times had they fallen in the past 12 months? A *recurrent fall* was defined as any case with 2 or more falls in a 12-month period.

- 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
2. Were any of these falls non-accidental, i.e. with no apparent or obvious reason? An *unexplained fall* was defined as any case who answered yes to this question.
 3. Were they injured seriously enough to require medical treatment? An *injurious fall* was defined as any case who answered yes to this question.

Based on the above responses, cases of fall were subdivided into *simple* and *complex*:

- *Simple falls* – defined as any case of fall that was not also documented as *recurrent*, *unexplained* and/or *injurious*.
- *Complex falls* – defined as any case of fall that was also documented to be *recurrent*, *unexplained* and/or *injurious*.

Each variable (syncope, simple fall and complex fall) was coded 1 if the case criteria were met, and 0 if they were not, producing a dichotomous variable for analysis.

Predictor Variables

Syncope Falls Index (SYFI)

As per standard procedure (14), a FI can be constructed on any suitable database by considering a minimum of 30 deficits that need to satisfy the following criteria: (a) be associated with health status, and not simply attributes; (b) cover a range of systems; (c) not saturate too early; and (d) their prevalence must increase with age (excluding survivor effects). For SYFI, 40 deficits were identified a priori from the Wave 1 TILDA ISSDA database as potentially increasing the risk of syncope and falls (Table 1). Of the 40 SYFI items, 38 were collected via self-administered questionnaire; and handgrip strength and Mini Mental State Examination were collected by a trained interviewer.

As regards the grip strength measure in SYFI, cases were considered to have the deficit where grip strength was found to be in the range of the frailty phenotype (23) (see Table 1 for details). As regards medications, the list of anticholinergics, benzodiazepines and Z-drugs characterised in TILDA has been described elsewhere.(24) For the cognitive deficit, we selected an MMSE score of less than 24 points.(25)

1
2
3
4
5
6 Each deficit was coded 1 if it was present or 0 if it was not. In each case, the number of
7 deficits recorded as present was divided by the total number of deficits considered (i.e. 40)
8 to produce the SYFI. The continuous score was then divided into tertiles to define SYFI risk
9 groups: low risk (first tertile), intermediate risk (second tertile) and high risk (third tertile). In
10 addition, we operationalised the binary frailty cut-off recommended by Dent et al. (SYFY >
11 0.25).(26)
12
13
14
15
16
17
18

19 *Age and sex*

20
21 The age and sex of each participant were recorded at Wave 1. Age and sex have known
22 influences on syncope.(2, 27) The association between chronological age and the FI is
23 typically small because the FI captures variation in health status within individuals of the
24 same age.(28) Studies have shown that women consistently have higher frailty scores than
25 men but lower overall mortality. Thus, we corrected for these demographics in our analysis.
26
27
28
29
30
31
32
33

34 *Analysis*

35
36 All statistical analyses were carried out with IBM SPSS Statistics version 25 (IBM Corp.:
37 Armonk, NY). Descriptives were given as mean with standard deviation (SD), median with
38 interquartile range (IQR), or proportion (%). The association between SYFI and age was
39 measured with the 2-sided Spearman's correlation coefficient (r_s). Associations between
40 continuous and dichotomous variables were reported with the two-sided independent-
41 samples Mann-Whitney U Test, and associations between dichotomous variables with the
42 Chi-square test.
43
44
45
46
47
48
49

50 To investigate the ability of SYFI to predict risk of future *syncope*, *complex falls* and *simple*
51 *falls* after adjusting for age and sex, binary logistic regression was performed on each of the
52 dependent variables. For regression analysis purposes, SYFI was entered as an ordinal
53 predictor where Group 1 acted as control for calculation of the odds ratios (OR) of Group 2
54 and Group 3. 95% confidence intervals (CI) for ORs were calculated. The level of significance
55 was set at $P < 0.05$.
56
57
58
59
60

1
2
3 In the main analysis, participants for whom the SYFI was calculated at wave 1 but did not
4 have outcomes measured in subsequent waves were omitted from analysis. A sensitivity
5 analysis was performed by re-running the logistic regression models after performing
6 multiple imputation of missing outcome data.
7
8
9

10 11 12 13 14 Declaration of Sources of Funding

15
16
17 The four waves of TILDA were funded by the Irish Government, The Atlantic
18 Philanthropies and Irish Life, plc. Roman Romero-Ortuno is funded by a Grant from
19 Science Foundation Ireland under Grant number 18/FRL/6188. The funders had no
20 role in study design, data collection and analysis, preparation of the manuscript, or
21 decision to publish.
22
23
24
25
26
27
28
29

30 31 32 Ethical Approval

33
34 Ethical approval for each wave was obtained from the Faculty of Health Sciences Research
35 Ethics Committee at Trinity College Dublin, Ireland. All participants provided written
36 informed consent prior to inclusion in the study.
37
38
39
40

41 42 43 Conflicts of Interest

44
45 The authors declare they have no conflicts of interest.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results

Of 8504 Wave 1 participants, 3499 were aged 65 or more years. Among the latter, the mean (SD) age was 72.6 (5.2) years, 52.5% were women, and all had SYFI information. In $N = 3499$, the r_s between age and SYFI was 0.214 ($P < 0.001$). Figure 1 shows the typical gamma distribution of SYFI across Wave 1 participants ($N = 3499$). At baseline, SYFI tertiles were: Group 1 or low-risk (SYFI < 0.05), Group 2 or intermediate-risk (SYFI $0.05 - 0.13$) and Group 3 or high-risk (SYFI > 0.13). 126 (3.6%) Wave 1 participants had a SYFI of more than 0.25.

By Wave 4, 592 participants (16.9% of Wave 1 sample), did not have information on outcomes. Table 2 shows a comparison of Wave 1 characteristics between participants with and without data for outcomes at Wave 4. The median age of participants without Wave 4 outcomes was three years older, but there were no significant differences in sex or median SYFI.

Of the 2907 with outcome data, 185 (6.4%) reported new syncope, 1077 (37.0%) complex falls, and 218 (7.5%) simple falls. By frailty status (SYFI > 0.25), 9.2% ($N = 9$) of the frail versus 6.3% ($N = 176$) of the non-frail had new syncope (Chi-squared test; $P = 0.245$); the proportions for complex falls were 62.2% ($N = 61$) and 36.2% ($N = 1016$) ($P < 0.001$); and for simple falls, 6.1% ($N = 6$) and 7.5% ($N = 212$) ($P = 0.599$), respectively.

The results of the multivariate binary logistic regression models (Table 3) showed that baseline SYFI had a statistically significant independent association with events of syncope and complex falls between Wave 1 and Wave 4, with incremental effects by tertiles. In contrast, SYFI did not show any statistically significant association with future simple falls. Age and sex were not shown to be significant predictors of simple falls or syncope. A small-magnitude association was observed with age and complex falls. Female sex appeared to be a significant independent predictor of complex falls (Table 3).

Appendix 1 shows a sensitivity analysis of the regression models utilising SYFI dichotomised as per frailty cut-off (> 0.25). Results showed that SYFI > 0.25 was an independent predictor of complex falls, but the syncope prediction lost significance. Appendix 2 shows another sensitivity analysis with imputation of missing data. Using age as a predictor for the imputation, we imputed values of missing outcomes data by using the SPSS automatic multiple imputation procedure. Results were unchanged compared to the main analysis.

Discussion

Using data from a large, community-based, nationwide study, we were able to compile an index (SYFI) that predicts, independently of age and sex, risk of future syncope and complex falls in community-dwelling older adults.

Syncope in older people is known to be complex and multifactorial. Advancing age, polypharmacy and multiple comorbidities are associated with physiological impairments in cardiovascular autoregulatory mechanisms, cerebral blood flow and neurohumoral stability.(29, 30) Frailty, cognitive decline, and reductions in functional status associated with ageing also contribute to the risk of syncope.(3) SYFI was designed to reflect this complex interplay between contributing factors, made up of a heterogenous group of 40 potential deficits that allows calculation of cumulative risk.

One of the clinical challenges in addressing syncope and complex falls is the underreporting and mislabelling of syncopal events. Atypical syncope presentations are common in older adults, the most prevalent being unexplained falls.(12, 31). We found that the risk of complex falls similarly rose for patients as they moved up SYFI risk groups. This suggests, as often clinically suspected, that complex falls in older people may have an underlying syncopal aetiology.(32) Female sex was in significant association with complex falls and this is also in keeping with known sex differences in syncope and falls epidemiology.(27)

Contrastingly, we found that SYFI was not predictive of future simple falls. The overlap in cardiovascular risk factors for both syncope and non-accidental falls has previously been described in this cohort.(32) In clinical practice, verifying whether non-accidental falls in older people have underlying syncopal aetiology has remained a challenge.(12, 33) General consensus supports the treatment of unwitnessed or unexplained falls as possible syncope (1, 5) but this is still underappreciated in non-specialist settings. Significant overlap in risk

1
2
3 profiles was identified in our study which further supports the mandate that unexplained,
4 recurrent and/or injurious falls in older people be considered as potential syncope.
5
6 Therefore, in patients with a high SYFI risk, it would be prudent to refer falls presentations
7 for further investigation. This could be particularly helpful in busy Emergency Departments
8 or primary care settings, where access to currently validated investigations for syncope,
9 such as tilt-table testing, is limited.
10
11
12
13
14
15
16

17 One of the advantages of the SYFI is that, although it incorporates a wide range of deficits,
18 the vast majority were measured using a self-administered questionnaire, which could easily
19 be recreated in a clinical setting. In addition, the SYFI only requires the measurement of grip
20 strength (handheld dynamometer), BMI (weight and height), and administration of the
21 MMSE. This lends itself to implementation as a risk assessment tool in non-specialist clinical
22 settings. Early detection and intervention for those at risk of syncope or complex falls, e.g.
23 referral to a syncope unit, comprehensive geriatric assessment and/or cardiac monitoring
24 devices, have been associated with favourable outcomes.(3, 11, 34) The SYFI could hence be
25 used to identify those who might benefit from these interventions.
26
27
28
29
30
31
32
33
34
35

36 A limitation of the SYFI is its strong reliance on self-report. However self-reported frailty
37 indexes have been validated elsewhere.(35) In our study, we found that 4.9% of the Wave 1
38 sample had cognitive impairment (MMSE < 24). This is likely to be of higher in real clinical
39 settings. Even without cognitive impairment, we cannot exclude recall bias as a potential
40 limitation of our study.
41
42
43
44
45
46
47

48 Another limitation is that only 126 Wave 1 participants had a SYFI of more than 0.25, which
49 is reflective of a generally healthy community-based sample with low proportions of severe
50 frailty. Results showed that SYFI > 0.25 was an independent predictor of complex falls, but
51 the syncope prediction lost significance probably due to the low numbers involved.
52
53
54
55
56
57
58
59
60

Conclusion

Using data from the TILDA cohort spanning 6 years, we were able to conceive an adapted frailty index that significantly predicts future risk of syncope and complex falls in those aged 65 years and over. Further assessment of the SYFI in clinical settings is recommended.

Tables and Figures

Table 1: SYFI deficits and their definitions.

	Deficit	Measure
1.	<i>Polypharmacy</i>	Taking five or more regular medications (excluding supplements)
2.	<i>Anti-hypertensive treatment</i>	Taking one or more anti-hypertensive medications
3.	<i>Anti-cholinergic use</i>	Taking one or more anti-cholinergic medications regularly (see methods for details)
4.	<i>Benzodiazepine use</i>	Taking one or more benzodiazepines regularly (see methods for details)
5.	<i>Z-drug use</i>	Taking one or more Z-drugs regularly (see methods for details)
6.	<i>Anti-depressant use</i>	Taking one or more antidepressants regularly
7.	<i>Weight loss</i>	Self-reported weight loss of more than 4.5kg or 10lb in the past 12 months
8.	<i>Poor eyesight</i>	Self-rated eyesight as poor (with or without corrective lenses)
9.	<i>Poor hearing</i>	Self-rated hearing as poor (with or without a hearing aid)
10.	<i>Poor smell</i>	Self-rated sense of smell as poor
11.	<i>Poor hearing</i>	Self-rated sense of taste as poor
12.	<i>Reduced Grip Strength</i>	Two measures of handgrip strength were taken using a hydraulic hand dynamometer (Baseline, Fabrication Enterprises, Inc., White Plains, NY) from the dominant hand, and the mean of these readings was calculated.

		<p><u>Cut-offs:</u></p> <p><i>Men:</i> 20.5kg – body mass index (BMI) < 24; 21.5kg – BMI 24–26; 23kg – BMI >26;</p> <p><i>Women:</i> 11.5kg – BMI < 23; 13kg – BMI > 23</p>
13.	<i>Hypertension</i>	Told by a doctor they had “high blood pressure or hypertension”
14.	<i>Angina</i>	Told by a doctor they had “angina”
15.	<i>Myocardial Infarction (MI)</i>	Told by a doctor they had “a heart attack (including MI)”
16.	<i>Congestive Cardiac Failure</i>	Told by a doctor they had “congestive heart failure”
17.	<i>Heart Murmur</i>	Told by a doctor they had “a heart murmur”
18.	<i>Arrhythmia</i>	Told by a doctor they had an abnormal heart rhythm
19.	<i>Stroke</i>	Told by a doctor they had “a stroke”
20.	<i>Transient Ischaemic Attack (TIA)</i>	Told by a doctor they had “a mini-stroke or a TIA”
21.	<i>Diabetes Mellitus</i>	Told by a doctor they had “diabetes”
22.	<i>Diabetic Ulcers</i>	Told by a doctor they had “leg ulcers”
23.	<i>Proteinuria</i>	Told by a doctor they had “protein in the urine”
24.	<i>Peripheral neuropathy</i>	Told by a doctor they had a “lack of feeling or tingling pain in the legs and feet due to nerve damage”
25.	<i>Diabetic retinopathy</i>	Told by a doctor that they had “damage to the back of the eye” related to diabetes
26.	<i>Diabetic nephropathy</i>	Told by a doctor they had “damage to the kidneys” related to diabetes
27.	<i>Chronic Obstructive Airway Disease</i>	Told by a doctor that they had “chronic lung disease such as chronic bronchitis or emphysema”

1		
2		
3		
4		
5		
6		
7		
8		
9		
10		
11		
12		
13		
14		
15		
16		
17		
18		
19		
20		
21		
22		
23		
24		
25		
26		
27		
28		
29		
30		
31		
32		
33		
34		
35		
36		
37		
38		
39		
40		
41		
42		
43		
44		
45		
46		
47		
48		
49		
50		
51		
52		
53		
54		
55		
56		
57		
58		
59		
60		
	28.	<i>Asthma</i>
		Told by a doctor that they had “asthma”
	29.	<i>Arthritis</i>
		Told by a doctor that they had “arthritis (including osteoarthritis)”
	30.	<i>Osteoporosis</i>
		Told by a doctor that they had “osteoporosis. or sometimes called thin or brittle bones”
	31.	<i>Malignancy</i>
		Told by a doctor that they had “cancer or a malignant tumour (including leukaemia or lymphoma)”
	32.	<i>Mental Health Disorder</i>
		Told by a doctor that they had “any emotional, nervous or psychiatric problem, such as depression or anxiety”
	33.	<i>Alcohol or Substance Abuse</i>
		Told by a doctor that they had “alcohol or substance abuse”
	34.	<i>Gastric Ulcers</i>
		Told by a doctor that they had “stomach ulcers”
	35.	<i>Varicose Ulcers</i>
		Told by a doctor that they had “varicose ulcers or an ulcer due to varicose veins”
	36.	<i>Liver Cirrhosis</i>
		Told by a doctor that they had “cirrhosis or serious liver damage”
	37.	<i>Cognitive impairment</i>
		Score less than 24/30 in the Mini-Mental State Examination (MMSE) test
	38.	<i>Urinary Incontinence</i>
		Self-reported “loss of urine beyond their control” in past 12 months
	39.	<i>Unsteadiness on Standing</i>
		Self-reported feeling unsteady on standing
	40.	<i>Unsteadiness on Getting Up from a Chair</i>
		Self-reported feeling unsteady on getting up from a chair

Table 2: Comparison of Wave 1 characteristics between participants with (N = 2907) and without (N = 592) data for outcomes at Wave 4.


	Outcome data available (N = 2907)	Outcome data missing (N = 592)	P value
Median age, years (IQR)	72 (9)	75 (10)	<0.001 [#]
Female sex (%)	52.6	50.0	0.870 [^]
Mean SYFI (SD)	0.08 (0.10)	0.10 (0.10)	0.440 [#]

IQR: interquartile range; [#] Independent-Samples Mann-Whitney U Test; [^] Chi-square test.

Table 3: Baseline SYFI groups (tertiles) as predictor of outcomes: results of the binary logistic regression models controlling for age and sex.

<i>Predictor</i>	<i>Odds Ratio [95% Confidence Interval]</i>	<i>P value</i>
Future syncope		
SYFI Group 1	<i>Control</i>	<i>Control</i>
SYFI Group 2	1.51 [1.03 – 2.21]	0.034
SYFI Group 3	1.88 [1.26 – 2.80]	0.002
Age	1.02 [0.99 – 1.05]	0.218
Female sex	1.06 [0.78 – 1.43]	0.724
Future complex fall		
SYFI Group 1	<i>Control</i>	<i>Control</i>
SYFI Group 2	1.34 [1.12 – 1.61]	0.002
SYFI Group 3	2.22 [1.82 – 2.72]	<0.001
Age	1.01 [1.00 – 1.03]	0.033
Female sex	1.51 [1.29 – 1.76]	<0.001
Future simple fall		
SYFI Group 1	<i>Control</i>	<i>Control</i>
SYFI Group 2	1.09 [0.79 – 1.51]	0.596
SYFI Group 3	0.96 [0.66 – 1.39]	0.831
Age	1.00 [0.97 – 1.03]	0.961
Female sex	1.14 [0.86 – 1.50]	0.370

Figure 1. Distribution of SYFI across Wave 1 participants (N = 3499).



References

1. Brignole M, Moya A, de Lange FJ, Deharo JC, Elliott PM, Fanciulli A, et al. 2018 ESC Guidelines for the diagnosis and management of syncope. *Eur Heart J*. 2018;39(21):1883-948.
2. Goyal P, Maurer MS. Syncope in older adults. *J Geriatr Cardiol*. 2016;13(5):380-6.
3. Wong CW. Complexity of syncope in elderly people: a comprehensive geriatric approach. *Hong Kong Med J*. 2018;24(2):182-90.
4. Chen LY, Gersh BJ, Hodge DO, Wieling W, Hammill SC, Shen WK. Prevalence and clinical outcomes of patients with multiple potential causes of syncope. *Mayo Clin Proc*. 2003;78(4):414-20.
5. Kenny RA. Syncope in the elderly: diagnosis, evaluation, and treatment. *J Cardiovasc Electrophysiol*. 2003;14(9 Suppl):S74-7.
6. de Ruyter SC, de Jonghe JFM, Germans T, Ruyter JH, Jansen RWMM. Cognitive Impairment Is Very Common in Elderly Patients With Syncope and Unexplained Falls. *J Am Med Dir Assoc*. 2017;18(5):409-13.
7. Campbell AJ, Buchner DM. Unstable disability and the fluctuations of frailty. *Age Ageing*. 1997;26(4):315-8.
8. Linzer M, Pontinen M, Gold DT, Divine GW, Felder A, Brooks WB. Impairment of physical and psychosocial function in recurrent syncope. *J Clin Epidemiol*. 1991;44(10):1037-43.
9. Kenny RA, Bhangu J, King-Kallimanis BL. Epidemiology of syncope/collapse in younger and older Western patient populations. *Prog Cardiovasc Dis*. 2013;55(4):357-63.
10. McCarthy K, Ward M, Romero Ortuño R, Kenny RA. Syncope, Fear of Falling and Quality of Life Among Older Adults: Findings From the Irish Longitudinal Study on Aging (TILDA). *Front Cardiovasc Med*. 2020;7:7.
11. Solbiati M, Costantino G, Casazza G, Dipaola F, Galli A, Furlan R, et al. Implantable loop recorder versus conventional diagnostic workup for unexplained recurrent syncope. *Cochrane Database Syst Rev*. 2016;4:CD011637.
12. O' Brien H, Anne Kenny R. Syncope in the Elderly. *Eur Cardiol*. 2014;9(1):28-36.
13. Mitnitski AB, Mogilner AJ, Rockwood K. Accumulation of deficits as a proxy measure of aging. *ScientificWorldJournal*. 2001;1:323-36.
14. Searle SD, Mitnitski A, Gahbauer EA, Gill TM, Rockwood K. A standard procedure for creating a frailty index. *BMC Geriatr*. 2008;8:24.
15. Kojima G, Iliffe S, Walters K. Frailty index as a predictor of mortality: a systematic review and meta-analysis. *Age Ageing*. 2018;47(2):193-200.
16. Legge A, Kirkland S, Rockwood K, Andreou P, Bae SC, Gordon C, et al. Construction of a Frailty Index as a Novel Health Measure in Systemic Lupus Erythematosus. *J Rheumatol*. 2020;47(1):72-81.
17. TILDA. The Irish Longitudinal study on Ageing (TILDA) Wave 1, 2009-2011. 1.9 ed. Irish Social Science Data Archive2019.
18. TILDA. The Irish Longitudinal study on Ageing (TILDA) Wave 2, 2012-2013. 2.4 ed. Irish Social Science Data Archive2019.
19. TILDA. The Irish Longitudinal study on Ageing (TILDA) Wave 3, 2014-2015. Irish Social Sciences Data Archive2019.
20. TILDA. The Irish Longitudinal study on Ageing (TILDA) Wave 4, 2016. 4.1 ed. Irish Social Sciences Data Archive2020.

21. Kearney PM, Cronin H, O'Regan C, Kamiya Y, Savva GM, Whelan B, et al. Cohort profile: the Irish Longitudinal Study on Ageing. *Int J Epidemiol*. 2011;40(4):877-84.
22. Donoghue OA, McGarrigle CA, Foley M, Fagan A, Meaney J, Kenny RA. Cohort Profile Update: The Irish Longitudinal Study on Ageing (TILDA). *Int J Epidemiol*. 2018;47(5):1398-l.
23. Savva GM, Donoghue OA, Horgan F, O'Regan C, Cronin H, Kenny RA. Using timed up-and-go to identify frail members of the older population. *J Gerontol A Biol Sci Med Sci*. 2013;68(4):441-6.
24. Moriarty F, Savva GM, Grossi CM, Bennett K, Fox C, Maidment I, et al. Cognitive decline associated with anticholinergics, benzodiazepines and Z-drugs: Findings from The Irish Longitudinal Study on Ageing (TILDA). *Br J Clin Pharmacol*. 2020.
25. Creavin ST, Wisniewski S, Noel-Storr AH, Trevelyan CM, Hampton T, Rayment D, et al. Mini-Mental State Examination (MMSE) for the detection of dementia in clinically unevaluated people aged 65 and over in community and primary care populations. *Cochrane Database Syst Rev*. 2016(1):CD011145.
26. Dent E, Kowal P, Hoogendijk EO. Frailty measurement in research and clinical practice: A review. *Eur J Intern Med*. 2016;31:3-10.
27. Park J, Jang SY, Yim HR, On YK, Huh J, Shin DH, et al. Gender difference in patients with recurrent neurally mediated syncope. *Yonsei Med J*. 2010;51(4):499-503.
28. Pérez-Zepeda MU, Godin J, Armstrong JJ, Andrew MK, Mitnitski A, Kirkland S, et al. Frailty among middle-aged and older Canadians: population norms for the frailty index using the Canadian Longitudinal Study on Aging. *Age Ageing*. 2020.
29. Güldner S, Langada V, Popp S, Heppner HJ, Mang H, Christ M. Patients with syncope in a German emergency department: description of patients and processes. *Dtsch Arztebl Int*. 2012;109(4):58-65.
30. Joy PS, Kumar G, Olshansky B. Syncope: Outcomes and Conditions Associated with Hospitalization. *Am J Med*. 2017;130(6):699-706.e6.
31. de Rooter SC, Wold JFH, Germans T, Rooter JH, Jansen RWMM. Multiple causes of syncope in the elderly: diagnostic outcomes of a Dutch multidisciplinary syncope pathway. *Europace*. 2018;20(5):867-72.
32. Cronin H, Kenny RA. Cardiac causes for falls and their treatment. *Clin Geriatr Med*. 2010;26(4):539-67.
33. Shaw FE, Kenny RA. The overlap between syncope and falls in the elderly. *Postgrad Med J*. 1997;73(864):635-9.
34. Kenny RA, Brignole M, Dan GA, Deharo JC, van Dijk JG, Doherty C, et al. Syncope Unit: rationale and requirement--the European Heart Rhythm Association position statement endorsed by the Heart Rhythm Society. *Europace*. 2015;17(9):1325-40.
35. Theou O, O'Connell MD, King-Kallimanis BL, O'Halloran AM, Rockwood K, Kenny RA. Measuring frailty using self-report and test-based health measures. *Age Ageing*. 2015;44(3):471-7.


Appendix 1: Baseline SYFI groups (Frail: SYFI > 0.25 vs Non-frail) as predictor of outcomes: results of the binary logistic regression models controlling for age and sex.

<i>Predictor</i>	Odds Ratio [95% Confidence Interval]	P value
Future syncope		
SYFI Frail	1.46 [0.72 – 2.95]	0.295
Age	1.03 [1.00 – 1.06]	0.066
Female sex	1.08 [0.80 – 1.45]	0.623
Future complex fall		
SYFI Frail	2.82 [1.86 – 4.29]	<0.001
Age	1.03 [1.01 – 1.04]	<0.001
Female sex	1.54 [1.32 – 1.79]	<0.001
Future simple fall		
SYFI Frail	0.80 [0.34 – 1.84]	0.593
Age	1.00 [0.97 – 1.03]	0.967
Female sex	1.14 [0.86 – 1.50]	0.370

Appendix 2: Baseline SYFI groups (tertiles) as predictor of outcomes: results of the binary logistic regression models controlling for age and sex, with multiple imputation of missing outcome data.

<i>Predictor</i>	<i>Odds Ratio [95% Confidence Interval]</i>	<i>P value</i>
Future syncope		
SYFI Group 1	<i>Control</i>	<i>Control</i>
SYFI Group 2	1.54 [1.05 – 2.25]	0.026
SYFI Group 3	1.89 [1.26 – 2.82]	0.002
Age	1.02 [0.99 – 1.05]	0.261
Female sex	1.04 [0.77 – 1.41]	0.781
Future complex fall		
SYFI Group 1	<i>Control</i>	<i>Control</i>
SYFI Group 2	1.35 [1.12 – 1.62]	0.001
SYFI Group 3	2.22 [1.82 – 2.71]	<0.001
Age	1.02 [1.00 – 1.03]	0.031
Female sex	1.51 [1.29 – 1.76]	<0.001
Future simple fall		
SYFI Group 1	<i>Control</i>	<i>Control</i>
SYFI Group 2	1.09 [0.79 – 1.51]	0.584
SYFI Group 3	0.96 [0.66 – 1.40]	0.843
Age	1.00 [0.97 – 1.03]	0.971
Female sex	1.14 [0.86 – 1.50]	0.369

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Distribution of SYFI across Wave 1 participants (N = 3499)

Downloaded from <https://academic.oup.com/qjmed/advance-article/doi/10.1093/qjmed/hcab141/6278904> by Trinity College Dublin user on 20 May 2021